

January 2015

STRESS GRANULES MODULATE SYK TO CAUSE MICROGLIAL DYSFUNCTION IN ALZHEIMER'S DISEASE

Soumitra Ghosh
Purdue University

Follow this and additional works at: https://docs.lib.purdue.edu/open_access_dissertations

Recommended Citation

Ghosh, Soumitra, "STRESS GRANULES MODULATE SYK TO CAUSE MICROGLIAL DYSFUNCTION IN ALZHEIMER'S DISEASE" (2015). *Open Access Dissertations*. 1412.
https://docs.lib.purdue.edu/open_access_dissertations/1412

This document has been made available through Purdue e-Pubs, a service of the Purdue University Libraries. Please contact epubs@purdue.edu for additional information.

PURDUE UNIVERSITY
GRADUATE SCHOOL
Thesis/Dissertation Acceptance

This is to certify that the thesis/dissertation prepared

By Soumitra Ghosh

Entitled

Stress Granules modulate SYK to cause microglial dysfunction in Alzheimer's disease

For the degree of Doctor of Philosophy



Is approved by the final examining committee:

Robert L. Geahlen

Chair

Jean Christophe Rochet

Riyi Shi

Val J. Watts

To the best of my knowledge and as understood by the student in the Thesis/Dissertation Agreement, Publication Delay, and Certification Disclaimer (Graduate School Form 32), this thesis/dissertation adheres to the provisions of Purdue University's "Policy of Integrity in Research" and the use of copyright material.

Approved by Major Professor(s): Robert L. Geahlen

Approved by: Jean Christophe Rochet

Head of the Departmental Graduate Program

8/11/2015

Date

STRESS GRANULES MODULATE SYK TO CAUSE MICROGLIAL
DYSFUNCTION IN ALZHEIMER'S DISEASE

A Dissertation

Submitted to the Faculty

of

Purdue University

by

Soumitra Ghosh

In Partial Fulfillment of the

Requirements for the Degree

of

Doctor of Philosophy

December 2015

Purdue University

West Lafayette, Indiana

To my parents and my best friend,
Anukana
who never stopped in believing and supporting me.

ACKNOWLEDGEMENTS

My imagination about USA was mainly influenced by movies; and my expectations were high. Though the first appearance of the town did not match my imagination, today I feel emotionally connected to Purdue University and West Lafayette. Purdue will always be there in my heart and play a major role in shaping my future. The excellent educational opportunity Purdue provided me throughout my PhD work far exceeds my expectations. There are many people who helped and encouraged me to achieve success and fight failures during my graduate work. It would be my great pleasure to thank and acknowledge them for their immense help towards successful completion of my graduate studies.

First I would like to thank my advisor, Dr. Robert Geahlen, for believing in me and accepting me as a graduate student to pursue exciting science in his laboratory. He was always there for me and supported me in every possible way to pursue science without boundaries. His valuable guidance and freedom to pursue science helped me to come up with such an interesting study. He taught me to be an independent researcher and critically think about any scientific problem. The patience he showed in correcting my documents through my graduate studies is worth mentioning. The long discussions we had about scientific problems and non-scientific topics would be cherished forever.

I would also like to thank Dr. Chris Rochet, Dr. Val Watts and Dr. Riyi Shi for taking time to serve on my advisory committee and keeping me on track through my graduate studies. Their valuable inputs and feedback helped me to make my research more interesting and challenging. I would also like to thank Dr. Watts for helping me as a mentor in graduate school. Dr. Donald Ready deserves special thanks for training me on

confocal microscopy and constantly taking care of the microscope so that we could continue our experiments throughout the year.

I would sincerely like to thank my lab mates with whom I have spent most of my time in the last few years. I have shared my frustrations and successes with them all the time. I would like to thank Shana and Mariya for spending time with me at breaks, debating on various topics, listening to me and helping me take care of my cells. I would also like to thank Mariya for accompanying me in scavenging free food/ goodies throughout the campus. I would like to thank Wen for providing me cell lines and sequencing the mouse genome for my studies. I have learned a lot from them that I can implement later in life. Finally I would like to thank my parents and my beloved Anukana for constantly believing in me and supporting me. Though my parents are unaware of what I do, they always stood by me and helped me in every possible way. Without their support, I would not be able to pursue this dream.

Most importantly, Anukana deserves special thanks for her contributions in my life as well as my studies. Her love, affection and support kept me going. She was the one who constantly made me feel better when I was stressed about my research and life. She would listen to me continuously about my experimental successes and failures. She is the best friend and partner I can ask for and deeply appreciate all that she has done for me.

TABLE OF CONTENTS

	Page
LIST OF TABLES	viii
LIST OF FIGURES	ix
LIST OF ABBREVIATIONS	xii
ABSTRACT	xvi
CHAPTER ONE: INTRODUCTION	1
1.1 Introduction	1
1.2 Alzheimer's disease and its causes	4
1.3 Amyloid beta and Tau pathology	4
1.4 Oxidative stress in AD	7
1.5 Genes, proteins and kinases involved in AD	7
1.6 Neuroinflammation in AD	13
1.7 Astrocytes-Activation and Inflammation	14
1.8 Microglia-Macrophages of the Brain	15
1.9 Various functions of Microglia	18
1.9.1 Phagocytosis of debris and foreign bodies	18
1.9.2 Engulfment of apoptotic cells	19
1.9.3 Participation in neurogenesis and brain wiring	20
1.9.4 Maintenance of Synaptic pruning	20
1.10 Microglial cells in Alzheimer's disease	21
1.11 SYK kinase and its activation	24
1.12 Receptor based immune signaling of SYK	27
1.12.1 Syk in B cell signaling	27
1.12.2 Syk in mast cell signaling	28
1.12.3 Syk in macrophage signaling	29
1.12.4 Syk in neutrophil signaling	31
1.13 Syk in brain	35
1.13.1 Syk in neuron	35
1.13.2 Syk in microglia	36

	Page
1.14 Stress granules and their components	39
1.14.1 SG assembly and disassembly	39
1.14.2 SG components	42
1.14.3 SG function	43
1.15 Stress granules in different neurological disease	46
1.15.1 SGs in Alzheimer's disease	46
1.15.2 SGs in Amyotrophic Lateral Sclerosis	46
1.15.3 SG in spinal muscular atrophy	47
 CHAPTER TWO: MATERIAL AND METHODS	 48
2.1 Cells and cell lines	48
2.2 Plasmids and Lentiviral vectors	49
2.3 Primary cell culture and Animal models	50
2.4 Stress granule formation	51
2.5 Western blotting and Immunoprecipitation assays	52
2.6 Microglial cell functional assays	53
2.7 Microglial-neuron co-culture and Annexin V assay	56
2.8 Human AD clinical samples	56
2.9 Antibodies	57
 CHAPTER THREE: RESULTS	 58
3.1 SYK is highly expressed in BV-2 and N9 microglia cell	58
3.2 Low dose of sodium arsenite (SA) promotes SG formation in microglial cells	60
3.3 Soluble amyloid beta induces SG formation in cells	62
3.4 SYK and G3BP1 co-localizes with other SG core in BV-2 and N9 cells	64
3.5 SYK associates with other SG core components in BV-2 cells	68
3.6 SYK is active when present in SGs	70
3.7 SYK regulates SG formation in MG under SA and A β treatment	73
3.8 Knockdown of SYK prevents SGs formation in MG under SA and under SA and A β treatment	75
3.9 SYK promotes SG formation in primary MG	78
3.10 Prolonged chronic stress induces large and irreversible SGs in N9 cells	83
3.11 Chronic stress induction lead to SYK-dependent generation of intracellular and extracellular ROS	85
3.12 Chronic stress induction lead to SYK-dependent generation of intracellular and extracellular RNS	87
3.13 Chronic stress induced MG cells mediated neuronal cell death through ROS and RNS secretion	89

	Page
3.14 Chronic stress induction in BV-2 and N9 cells impairs phagocytosis of pH sensitive E.Coli Rhodo particle.....	91
3.15 Chronic stress in SG in MG prevents phagocytosis of β -amyloid fibrils	93
3.16 SYK associated SGs are formed in brains of AD patients.....	95
3.17 IgG treatment can repair impaired phagocytosis after chronic stress induction.....	100
CHAPTER FOUR: DISCUSSION	105
CHAPTER FIVE: FUTURE DIRECTIONS	112
LIST OF REFERENCES	117
VITA.....	149

LIST OF TABLES

Table	Page
1.1 Characteristics of Alzheimer's disease patients used to measure SYK, G3BP1 and phosphotyrosine (pTYR) co-localized in SGs	97

LIST OF FIGURES

Figure	Page
1.1 Schematic diagram of Alzheimer's disease hallmarks	6
1.2 Schematic chart of Alzheimer's disease causative factors	12
1.3 Schematic diagram of microglial cell in AD	17
1.4 Schematic diagram of SYK structure	25
1.5 Schematic diagram of adaptor mediated SYK-ITAM signaling.....	33
1.6 Schematic diagram of direct ITAM-SYK signaling	34
1.7 Schematic diagram of stress granules in neurodegeneration	46
3.1 SYK is highly expressed in N9 and BV-2 microglial cell	49
3.2 SA stimulation manifests SG formation in N9 and BV-2 cells	61
3.3 A β stimulation manifests SG formation in N9 and BV-2 cells	63
3.4 SYK co-localizes with pEIF2 α , TDP-43, PABP1 and TIA-1 in BV-2 and N9 cells...	65
3.5 G3BP1 co-localizes with pEIF2 α , TDP-43, PABP1 and TIA-1 in BV-2 cells	66
3.6 SG markers like SYK, G3BP1, TIA-1 and pEIF2 α expression increases in	soluble
fraction in BV-2 and N9 cells	67
3.7 SG markers G3BP1 and pEIF2 α associate with GFP-tagged SYK in SYK-EGFP expressing BV-2 cell.....	69
3.8 SYK is active when co-localized in SGs in N9 cells under treatment.....	71
3.9 SYK is active when co-localized in SGs in BV-2 cells under treatment.....	72
3.10 SG formation dynamics is affected by SYK.....	74
3.11 SYK knockdown prevents G3BP1 associated SG formation in BV-2 cells	76
3.12 SYK knockdown prevents SYK associated SG formation in BV-2 cells.....	77
3.13 Primary MG culture express IBA-1 and mice expressed SYK	79

Figure	Page
3.14 Adult primary MG from 20 month old mice generates large SGs upon stress induction compared to cells from 1 month old mice	80
3.15 Adult primary MG from 20 month old mice Syk ^{+/-} generate large SGs upon stress induction compared to cells from 1 month old Syk ^{+/-} mice	81
3.16 Quantitative comparison of SG formation in MG from young (1m) and old (20m), wild-type (Wt) and Syk haploinsufficient (Het) mice	82
3.17 Prolonged chronic stress stimulation manifests persistent and irreversible SG in N9 cells	84
3.18 Chronic stress stimulation leads to increased ROS generation in MG cells	86
3.19 Chronic stress stimulation leads to increased RNS generation by MG cell	88
3.20 Chronically stressed MG cells induce apoptosis of HT22 neuronal cells in MG-neuron co-culture	90
3.21 Chronic SA and A β stimulation and SYK inhibition impairs phagocytosis of E.coli Rhodo particles in BV cells	92
3.22 Chronic SA and A β stimulation impairs phagocytosis of FITC-A β fibrils in N9	94
3.23 SYK and G3BP1 positive granules were observed in human AD brain samples	97
3.24 SYK and G3BP1 positive granules co-localized in human AD brain samples	98
3.25 SYK positive granules co-localized with phosphotyrosine in human AD brain Samples	99
3.26 IgG treatment of chronically stressed N9 cells restores phagocytic activity	102
3.27 IgG treatment of chronically stressed N9 cells relocalizes SYK to the plasma membrane	103
3.28 IgG treatment of chronically stressed N9 cells partially relocalizes SYK from the SG insoluble fraction to the soluble fraction	104
4.1 Model of SYK associated stress granules formation in microglial cell	107
4.2 Graph art depicting stress granule formation in microglial cell upon A β	109

LIST OF ABBREVIATIONS

AMPK	5' adenosine monophosphate-activated protein kinase
ADAM10	A Disintegrin and metalloproteinase domain-containing protein 10
AP-1	Activator protein-1
ATP	Adenosine triphosphate
AD	Alzheimer's disease
APP	Amyloid precursor protein
ALS	Amyotrophic lateral sclerosis
APOE	Apolipoprotein E
ATG	Autophagy related protein
AUB	AU-binding proteins
BCR	B-cell receptor
A β	Beta-amyloid
β -CTF	Beta C-terminal fragments of APP
BDNF	Brain derived neurotrophic factor
BAI1	Brain-specific angiogenesis inhibitor 1
BRF1	B-Related Factor 1
BTK	Bruton's tyrosine kinase
CAMKK2	Calcium /calmodulin dependent kinase kinase 2
CaMKII	Calcium/ Calmodulin-dependent protein kinase II
CREB	cAMP response element binding protein
CNS	Central nervous system
CX3CL1	Chemokine fractalkine
JNK3	c-Jun N-terminal kinase 3
CSF1R	Colony stimulating factor 1 receptor
Cdk5	Cyclin dependent kinase 5
COX2	Cyclooxygenase 2
CNTF	Cytokine ciliary neurotrophic factor
CPEB	Cytoplasmic polyadenylation element binding protein
DAMPs	Damage associated molecular patterns
DAP12	DNAX-Activation Protein 12
Elk-1	E26-like kinase 1
EAD	Early-onset AD
EPHB2	Ephrin receptor B2
EMPs	Erythromyeloid progenitors
eIF2 α	Eukaryotic translation initiation factor 2-alpha
ERK	Extracellular-signal-regulated kinases

FASL	Fas Ligand
FcεR1	Fc Epsilon receptor 1
FBP	Folate-binding protein
FMRP	Fragile X mental retardation protein
FXR1	Fragile X mental retardation related protein 1
FLD	Frontotemporal lobar degeneration
FTLD	Frontotemporal lobar degeneration
FUS	Fused in Sarcoma
GCN2	General control nonderepressible 2
GWAS	Genome wide association studies
GPX	Glutathione peroxidase
GST	Glutathione S-transferase
GSH	Glutathione
GW182	Glycine-tryptophan 182
GSK-3β	Glycogen synthase kinase 3 beta
GM-CSF	Granulocyte-macrophage colony-stimulating factor
GRB7	Growth factor receptor-bound protein 7
HRI	Heme-regulated eIF2α kinase
HDLS	Hereditary Diffuse Leukoencephalopathy with Spheroids
HDAC6	Histone deacetylase 6
HD	Huntington's disease
H ₂ O ₂	Hydrogen peroxide
OH ⁻	Hydroxyl anion
IgG	Immunoglobulin G
ITAM	Immunoreceptor tyrosine-based activation motif
ITIM	Immunoreceptor tyrosine-based inhibition motif
iNOS	Inducible nitric oxide synthase
IP5K	Ins(1,3,4,5,6)P5 2-kinase
IGF-1	Insulin-like growth factor 1
IFN _γ	Interferon gamma
IL-1β	Interleukin-1 beta
IL-1	Interleukin-1
LAD/ LOAD	Late-onset AD
LAT	Linker for activation of T cells
LPS	Lipopolysaccharide
MCP	Macrophage chemotactic protein
M-CSF	Macrophage-colony stimulating factor
mTOR	Mammalian target of rapamycin
MTK1	MAP kinase kinase kinase
mRNA	Messenger RNA
MG	Microglial cell
MCI	Mild cognitive impairment
MFG-E8	Milk fat globule epidermal growth factor 8
mmLDL	Minimally oxidized low-density lipoprotein
MAPK	Mitogen activated protein kinase

MEGF10	Multiple epidermal growth factor-like domains 10
MS	Multiple sclerosis
NGF	Nerve growth factor
NPC	Neuronal precursor cells
NT-3	Neurotrophin 3
NO	Nitric oxide
NMDA/ NR2B	N-methyl D-aspartate receptor
NSAID	Nonsteroidal anti-inflammatory drugs
NFAT	Nuclear factor of activated T cells
PNS	Peripheral nervous system
ONOO ⁻	Peroxynitrite
PS	Phosphatidylserine
PI3K	Phosphoinositide 3-kinase
PDK1	Phosphoinositide-dependent kinase-1
PLC γ	Phospholipase C γ
pTYR	Phosphotyrosine
PABP1	Poly(A)-binding protein 1
PSEN1/ PSEN2	Presenilin 1/ 2
PRNP	Prion protein
PGE ₂	Prostaglandin E ₂
PKA	Protein kinase A
PKR	Protein kinase R
PERK	Protein kinase RNA-like endoplasmic reticulum kinase
PP-2A	Protein phosphatase 2A
G3BP	Ras-GTPase-activating protein SH3-domain-binding protein
RNS	Reactive nitrogen species
ROS	Reactive oxygen species
RACK1	Receptor for activated C kinase 1
RAGE	Receptor for advanced glycation endproducts
RNP	Ribonucleoprotein
RAP55	RNA-associated protein 55
SLP76	SH2 domain leukocyte-specific phosphoprotein
SIGLECS	Sialic acid-binding immunoglobulin superfamily lectins
SIRP β 1	Signal regulatory protein- β 1
SA	Sodium arsenite
sA β	Soluble A β
SYK	Spleen tyrosine kinase
SH2	Src Homology 2
STAU1	Staufen1
SRC-3	Steroid receptor coactivator-3
SG/ SGs	Stress granules
O ₂ ⁻	Superoxide
SMN	Survival of motor neuron
TIA-1	T cell internal antigen-1
TIAR	T cell internal antigen-1-related

TDP-43	TAR DNA-binding protein 43
TIM4	T-cell immunoglobulin mucin receptor 4
TRAF2	TNF Receptor-Associated Factor 2
TLRs	Toll like receptors
TREM2	Triggering Receptor Expressed On Myeloid Cells 2
UPS	Ubiquitin-dependent proteasome system
TTP	Tristetraprolin
TNF- α	Tumor necrosis factor α
TYRO	TYRO Protein Tyrosine Kinase Binding Protein
VCP	Valosin-containing protein
ZBP1	Z-DNA Binding Protein 1
ZAP70	ζ -chain-associated protein kinase

ABSTRACT

Ghosh, Soumitra. PhD., Purdue University, December 2015. Stress Granules modulate Syk to cause Microglial Dysfunction in Alzheimer's disease. Major Professor: Robert L. Geahlen.

Microglial cells in the brains of Alzheimer's patients are recruited to amyloid beta ($A\beta$) plaques and exhibit an activated phenotype, but are defective for plaque removal by phagocytosis. To explore the molecular basis for these phenomena, I hypothesized that defects in the functions of the protein-tyrosine kinase SYK, which is important both for macrophage activation and phagocytosis, might underlie much of this pathology. Recent evidence from our lab indicates that SYK can associate with stress granules, ribonucleoprotein particles that form in stressed cells and contain inactive translation initiation complexes. I found that microglial cell lines and primary mouse brain microglia, when stressed by exposure to sodium arsenite or $A\beta(1-42)$ peptides or fibrils, form extensive stress granules to which the tyrosine kinase, SYK, is recruited. SYK enhances the formation of stress granules as evidenced by the inhibition of stress granule formation by small molecule inhibitors, knockdown of SYK expression by shRNA and SYK haploinsufficiency in mouse microglial cells. SYK is active within the resulting stress granules where it catalyzes the phosphorylation of stress granule-associated proteins on tyrosine. SYK-dependent stress granule

formation stimulates the production of reactive oxygen and nitrogen species. These are toxic to neuronal cells as demonstrated by a co-culture assay using stressed microglial cells and HT22 neuronal cells. The ability of microglial cells to phagocytose *E. coli* is blocked by SYK inhibitors. The sequestration of SYK into stress granules inhibits the ability of microglial cells to phagocytose either *E. coli* or A β fibrils. Microglial cells from aged mice are more susceptible to the formation of stress granules than are cells from young animals. Stress granules containing SYK and phosphotyrosine are prevalent in the brains of patients with severe Alzheimer's disease, suggesting that the sequestration of SYK into stress granules is part of the pathology of the disease. Phagocytic activity can be restored to stressed microglial cells by treatment with IgG independent of the epitope specificity, suggesting a mechanism to explain the therapeutic efficacy of intravenous IgG.

CHAPTER ONE: INTRODUCTION

1.1 **Introduction**

Microglial cells (MG) are the professional macrophages of the central nervous system. Among their many tasks is the removal from the brain of aggregates of β -amyloid ($A\beta$), which is formed from a proteolytic product of amyloid precursor protein (1). The persistent accumulation of β -amyloid plaques is a characteristic of Alzheimer's disease (AD), a serious neurodegenerative disorder affecting millions of patients worldwide. MG are attracted to sites of $A\beta$ deposition, which they recognize through a variety of cell surface receptors, and are capable of $A\beta$ removal through phagocytosis (2, 3). Impaired microglial activity, as exhibited by defective removal of $A\beta$ plaques, is particularly associated with the later stages of AD as plaques accumulate (4-6). These plaques still attract MG, but are refractory to phagocytosis. While MG can phagocytose $A\beta$ fibrils, they also become activated as a consequence of receptor engagement by $A\beta$ (7-9). The activation of MG is problematic as the resulting inflammatory response can damage neighboring neuronal cells (9). Compelling evidence for an important role of MG in AD comes, in part; from large scale studies of genes associated with AD that directly implicate inflammatory responses of MG as critical for AD pathology. Products of genes with identified associations with AD include Triggering Receptor Expressed on Myeloid

Cells 2 (TREM2), TYRO Protein Tyrosine Kinase Binding Protein (TYROBP) and CD33 (10-15). The myeloid receptor TREM2 functionally associates with TYROBP and promotes the internalization of bacteria and apoptotic neurons and recruits MG to A β plaques (16-19). Loss-of-function mutants promote inflammatory responses, decrease phagocytosis and predispose patients to AD (10). CD33 is dysregulated in MG in AD brain and is associated with reduced uptake of A β and increased numbers of activated, pro-inflammatory MG (20).

Many phagocytic receptors contain or are associated with proteins such as TYROBP that contain immunoreceptor tyrosine-based activation motifs (ITAMs) (21, 22). Receptor engagement initiates the phosphorylation of the two ITAM tyrosines leading to a physical interaction with the tyrosine kinase, SYK (23). Signaling through ITAM receptors is attenuated by receptors with immunoreceptor tyrosine-based inhibitory motifs (ITIMs) such as CD33 (24). Since SYK is essential for both phagocytosis in macrophages and for the receptor-mediated activation of inflammatory responses (25, 26), two events that are disrupted in MG from AD brains, it would appear to be a prime candidate for a critical mediator of AD pathology. An important question is how the function of SYK might be disrupted in aged or stressed MG.

MG in aged and AD brain fail to clear A β plaques, but display an activated, pro-inflammatory phenotype (4-6). How aging and other stresses affect MG to alter their homeostatic functions is unclear. Eukaryotic cells respond to external stresses via the formation of ribonucleoprotein (RNP) complexes scaffolded by RNA-binding proteins that self-associate. Examples of RNP aggregates include stress granules (SGs), which contain stalled initiation complexes and their associated mRNAs (27, 28). An association

of RNP aggregates with neurodegenerative diseases is well documented and mutations in RNA-binding proteins that promote self-assembly can be drivers of motor neuron diseases (29, 30). Similarly, defective clearance of SGs leads to the pathological accumulation of RNP particles and underlies the pathology of amyotrophic lateral sclerosis, Huntington's disease (HD), frontotemporal lobar degeneration (FLD) and AD (31-40).

Large scale proteomic screens from our laboratory have identified multiple SG-associated proteins as binding partners and substrates for SYK (41-43). SYK is recruited to SGs in some cells when exposed to sodium arsenite (SA) or proteasome inhibitors. In this study, I find that SYK also is recruited to SGs that form in MG, not only in response to sodium arsenite, but also in response to A β peptides or fibrils. Chronic stress leads to the formation of persistent SGs in MG in which active SYK is sequestered, leading to the constitutive production of reactive metabolites and the inhibition of phagocytic activity. Cytoplasmic aggregates containing SYK and phosphotyrosine are more pronounced in MG in the brains of patients with advanced AD. Based on this information, I propose a model in which the production of SGs and the corresponding misregulation of SYK directly contributes to AD pathology. Interestingly, the impaired phagocytic activity of stressed MG can be restored by the induced relocation of SYK from the SG to the plasma membrane by treating cells with Immunoglobulin G, suggesting a mechanism by which the administration of immunoglobulins to AD patients might retard disease progression.

1.2 Alzheimer's disease and its causes

Alzheimer's disease (AD) is one of the most prevalent neurodegenerative disorders in the world. According to recent statistics, five million Americans are living with AD in 2015, a number predicted to rise to sixteen million by 2050 (44). A substantial increase in the older population is also considered a major contributor to this alarming increase in AD patients. AD is categorized into two categories: early-onset AD (EAD) and late-stage AD (LAD). Mild-cognitive impairment (MCI) is a cognitive dysfunctional condition that precedes AD and contributes to its progression (45-48). Currently we do not have a method to detect AD in its early stages (49). The clinical symptom of AD develops much later than the initial onset of the disease; hence early detection is extremely difficult. The primary cause of dementia in AD is still unknown (50). The term AD was first coined in 1910 by Dr. Kraepelin following its discovery in 1906 by Dr. Alois Alzheimer. Considerable progress in understanding the complexity of the disease has not yet led to a successful therapeutic intervention. Hence we do not have a successful drug candidate to cure AD.

1.3 Amyloid beta and Tau pathology

The patients suffering from AD encounter reduced memory, cognition dysfunction and massive neuronal cell death over time. Post mortem examination of AD patients' brains have revealed amyloid beta ($A\beta$) plaques and tau fibrillary tangles as the two hallmarks of AD (10, 45, 46, 51-53). $A\beta$ aggregates into oligomers, fibers and plaques in the

diseased brain that hinder the normal functioning of brain in various ways (54-59). A β is amino acid peptide sequence of variable size (38, 40 and 42 amino acids) that is formed as result of cleavage of amyloid precursor protein (APP) by two enzymes: β -secretase and γ -secretase. It is considered that A β deposition is an initiating pathological event in AD. Over time A β undergoes different stages of aggregation, forming soluble oligomers, insoluble oligomers and insoluble fibrils (60-63). These A β deposits impair synaptic functioning resulting in chronic neurodegeneration ultimately leading to excessive neuronal cell death, cognitive impairment and severe dementia (64-67). Targeting A β deposition through drugs, antibodies and small molecule inhibitors have proven to be effective in alleviating the symptoms to a certain extent, but not completely (68-73).

Tau is another key protein involved in neurodegeneration. Tau is a microtubule-associated protein found in mature neurons where it stabilizes microtubules. In normal brain, tau is both O-GlcNAcylated and phosphorylated at low levels. In AD brain, hyperphosphorylation of tau leads to impairment of its normal biological activity. In neurodegenerative conditions, tau becomes N-glycosylated; and this glycosylation makes it more prone to phosphorylation by protein kinase A (PKA), glycogen synthase kinase 3 beta (GSK-3 β) and cyclin dependent kinase 5 (Cdk5). Additionally protein phosphatase A (PP-2A) activity is decreased, but calmodulin-dependent protein kinase II (CaMKII), PKA and mitogen activated protein kinase (MAPK) activities are increased in the CNS (69, 74-84). The hyperphosphorylation of tau leads to a breakdown of the microtubule network and subsequent formation of paired helical filaments which make up neurofibrillary tangles. Finally axonal transport is compromised resulting in retrograde degeneration and dementia (77-80, 85).

PATHOLOGICAL HALLMARKS OF ALZHEIMER'S DISEASE




Figure1.1 Schematic diagram of Alzheimer's disease hallmarks

In human AD brain, Amyloid beta protein aggregates to form amyloid plaques as shown in the figure. TAU protein also hyperphosphorylates to form neurofibrillary tangles as shown above.

Adapted from NIA-NIH webpage (Imaged by Colorado Hospital).

Scale=5 μ m

1.4 **Oxidative stress in AD**

Apart from A β and tau pathology, there are several other factors that contribute to AD pathology. Maintenance of cellular redox levels is crucial for the normal functioning of neurons. Studies on AD brains suggest excessive oxidative stress damages neurons and other supporting cells in the brain. Common oxidants such as hydrogen peroxide (H₂O₂), superoxide (O₂⁻) and peroxynitrite (ONOO⁻) are highly elevated in AD brains (74, 86-89). Additionally there is a decrease in antioxidants such as glutathione (GSH) and its supporting enzymes, glutathione peroxidase (GPX), and glutathione S-transferase (GST) (90). The increase in oxidative stress is mainly attributed to elevated oxidative phosphorylation in damaged mitochondria. AD progression impairs the normal efficiency of the mitochondria and leads to a feed-forward mechanism of superoxide production (91-93). Increase in ROS/ RNS promotes neuroinflammation through increase in cytokines, activation of apoptotic pathways and overexpression of transcription factors leading to neuronal cell death (85, 92, 94-100).

1.5 **Genes, proteins and kinases involved in AD**

AD is also classified into two categories, sporadic and familial, based on its causative origin. Our knowledge about the genes involved in AD mainly comes from studies involving familial AD cases. Studies for decades have revealed that several genes and proteins through elevated levels, increased activity and mutations contribute to AD progression.

Amyloid precursor protein (APP) is a type1 transmembrane protein that is highly expressed in neurons and astrocytes. Proteolytic processing of APP leads to amyloidogenic pathways (101, 102). APP is cleaved to generate A β peptides, soluble APP β and beta C-terminal fragments β -CTF (103, 104). APP on the cell surface is generally internalized allowing A β to be generated within the endocytic pathway and then released into the extracellular space (105). Recessive genetic mutations in APP such as A673V and E693 Δ are reported to cause early-onset AD (106). Mutations such as the Swedish mutation (KM670/671NL) increases plasma A β levels by altering β -secretase enzyme efficiency whereas the Dutch mutation (E693Q) occurs in the A β domain and results in excess A β aggregation (107, 108).

Presenilin 1 (PSEN1) and presenilin 2 (PSEN2) are integral membrane proteins that are important components of the γ -secretase complex, which is responsible for cleaving APP into A β fragments (109, 110). Their presence in the endoplasmic reticulum makes them an important component of protein processing. To date, 185 dominant mutations have been identified in *PSEN1* that account for 80% of early-onset familial AD and 13 dominant pathogenic mutations have been identified in *PSEN2* (111, 112).

A Disintegrin and metalloproteinase domain-containing protein 10 (ADAM10) is a metalloproteinase domain-containing protein that is considered a major α -secretase and is responsible for cleaving the APP ectodomain. ADAM10 mutants Q170H and R181G increase A β levels in vitro and in AD transgenic mice (113-116).

Apolipoprotein E (APOE) is a prime risk gene in AD. It is a pleiotropic glycoprotein that is highly expressed in liver, brain and macrophages. It regulates the mobilization and redistribution of cholesterol. APOE in the brain is required for neuronal

growth, nerve repair and regeneration, immune response and activation of lipolytic enzymes. APOE exists in three different isoforms: APOE ϵ 2, APOE ϵ 3 and APOE ϵ 4. APOE ϵ 4 has been reported to increase the risk of familial and sporadic early- and late-onset AD. The increase in APOE expression can increase plaque formation and accumulation (51, 106, 117-119).

Evidence suggests that cellular prion protein PRNP plays an important role in several neurodegenerative diseases including AD. It has a very high binding affinity for A β oligomers as confirmed in human AD brains. It binds to the tyrosine kinase FYN, which phosphorylates N-methyl D-aspartate receptor subtype 2B (NR2B) subunit and TAU on tyrosine. Interestingly, FYN phosphorylates TAU postsynaptically to induce A β toxicity mediated through the N-methyl-D-aspartate (NMDA) receptor (120-127).

Ephrin receptor B2 (EPHB2) is a tyrosine kinase that has a high affinity for A β oligomers and helps in regulating the synaptic localization and function of NMDA receptors. Studies have revealed that A β oligomers can bind EPHB2 and lead to its proteosomal degradation. This affects NMDA receptor signaling, resulting in defective long term potentiation (LTP) (66, 120, 128-130).

Evidence has suggested an impairment of calcium signaling in AD brains. A β induced calcium elevation activates the calcium-dependent phosphatase calcineurin, which in turn leads to the nuclear localization and increased transcriptional activity of nuclear factor of activated T cells (NFAT). GSK3 phosphorylates NFAT to keep it out of the nucleus which in turn increases Fas ligand (FASL) levels. Calcium /calmodulin dependent kinase kinase 2 (CAMKK2) is a serine /threonine kinase whose activity is directly dependent on calcium influx in the brain (131-138). The CAMKK2- 5' adenosine

monophosphate-activated protein kinase (AMPK) pathway mediates early synaptotoxic effects of A β oligomers through Tau phosphorylation (139, 140).

Mitochondrial damage is an early event observed in AD cell–culture and animal models. Centaurin- α 1 is an Arf GTPase-activating protein that promotes ADP ribosylation and is required for normal dendritic development. It interacts with RAS and Ras-E26-like kinase 1 (Elk-1) in AD brains. Mitochondrial permeability is directly regulated by centaurin- α 1 and Elk-1. Ras-Elk-1 pathway mediated mitochondrial damage leads to loss of dendritic spines and spine structural plasticity (115, 141-148).

5' adenosine monophosphate-activated protein kinase (AMPK) and mammalian target of rapamycin (mTOR) pathways are directly modulated by Ras and MAPK in neurons leading to neuronal cell death. mTOR is a serine/threonine kinase that plays a critical role in several cellular processes such as maintaining cell size, cell proliferation and autophagy through regulation of protein synthesis. Specifically in neurons, mTOR plays a role in synaptic plasticity, axon pathfinding and regeneration, dendrite arborization and spine morphology maintenance. In AD brains, mTOR signaling is unregulated in both mouse models and human samples. This manifests in an early cognitive decline and reduced autophagy that can be rescued by inhibition of mTOR by rapamycin.

Additionally, AMPK and MAPK4 further phosphorylate TAU on S262 in the microtubule-binding domain. This in turn promotes tau aggregation and hinders its clearance (149, 150).

Two other kinases that play an important role are Cdk5 and c-Jun N-terminal kinase 3 (JNK3). ER stress is a common phenomenon observed in AD. ER stress activates JNK3 that further phosphorylates APP. Thus increased JNK3 activity in AD

brain facilitates endocytosis of APP and further processing of A β (142, 151-160). Cyclin dependent Kinase (Cdk5) is a proline-directed serine/ threonine kinase that also plays a critical role in AD progression. In neurons, Cdk5 binds to its activator p35 or p39 to maintain its activation. Later in AD, Cdk5 is over activated by the p25 component of p35 and this leads to synaptic failure, which promotes neuronal cell death and loss of learning and memory. This overactivation leads to Tau aggregation, A β accumulation, activation of downstream transcription factors, and increased oxidative stress in the brain (90, 152, 161-166).

GSK3 β is modulated by Wnt signaling and insulin levels in the brain. GSK3 also reduces acetylcholine synthesis and promotes neuronal apoptosis. GSK3 β plays a pivotal role in AD signaling cascade leading to hyperphosphorylation of tau and increased A β production along with cerebral inflammation (162, 167, 168).




Figure 1.2 Schematic chart of Alzheimer's disease causative factors

In AD, both environmental and genetic factors contribute towards the manifestation of the disease. Environmental factors as mentioned above are considered the cause behind 80% AD cases. These factors could also lead to mutations in genes described in genetic factors. Genetic factors like Amyloid precursor protein interact with other genes and proteins that predispose individual to AD.

1.6 Neuroinflammation in AD

Neuroinflammation is a crucial event that contributes to AD pathogenesis and alters the neuronal compartment of the brain. Misfolded and aggregated protein complexes bind to pattern recognition receptors on microglia and astroglia triggering an innate immune response (131, 141, 169-174). This response results in the release of inflammatory factors that increase disease severity. High throughput screening studies have revealed that many genes that increase the risk for sporadic AD are actually encoded in glial cells. Other external factors like obesity, metabolic disorders, exposure to toxins, stress and high cholesterol all indicate interference with the immunological processes of the brain. Our ability to understand and control such immune reactions in the brain would be an important step to prevent or delay neurodegenerative diseases (69, 74, 87, 94, 141, 169-171, 175-178).

The innate immune response toward amyloid deposition and neurofibrillary tangle formation is considered a prime pathological event in AD. It has been observed that in AD brains amyloid plaques are always surrounded by activated microglial cells and astrocytes. The complement-mediated immune system has also been found to be activated in amyloid- and tau tangles-positive AD brains (78, 141, 179-181). Once the amyloid plaques or tangles are formed, the immune system mediates a local inflammatory response that directs the immune cells to the site of deposition and promotes secondary inflammation. Immune-based therapies have been designed to remove amyloid peptide aggregates from the brain of AD animal models. Several generic non-steroidal anti-inflammatory drugs have been shown to be slightly effective against

AD progression (100, 182-184). The nonsteroidal anti-inflammatory drug (NSAID) R-flurbiprofen was used in a recent Phase II clinical trial in patients affected with mild AD and showed a 62% decline in AD symptoms. These NSAID's mainly target cyclooxygenase 2 (COX2), which is found to be up regulated in AD brains and enhances neuroinflammation (182, 185-188). NSAIDs such as ibuprofen and indomethacin target cyclooxygenases nonselectively and suppress prostaglandin production by microglial cells during inflammatory reactions (189, 190). Few agents directly target β -secretase and APP-CTF- β thereby hindering proteolysis of APP and APP-CTF- β respectively. In the brain, the resident immune cells sense the alterations in the tissue through receptors for damage associated molecular patterns (DAMPs) (191). In this case, DAMPs are amyloid plaques, tau fibrillary tangles and misfolded proteins. The extreme inflammatory condition observed in AD is driven mainly by CNS-resident immune cells —microglia, perivascular myeloid cells and astrocytes.

1.7 Astrocytes- Activation and Inflammation

Astrocytes and microglia are the two major glial cell populations of the brain. They both communicate with neurons to maintain homeostasis and proper functioning of the brain. In addition to participating in the immune response, astrocytes recycle glutamate, and regulate CNS blood flow, blood brain barrier formation and maintenance, synaptogenesis, myelination, metabolism, and release of transmitters (64, 74, 97, 141, 192-195). They are the predominant cells in the mammalian brain. Their role has been documented in several neuro-developmental diseases such as Rett Syndrome and Fragile X Syndrome and in

neurodegenerative disorders such as multiple sclerosis (MS), Alzheimer's disease, amyotrophic lateral sclerosis (ALS) and HD (192, 196-199). Developmental studies have suggested that astrocytes develop by gliogenesis, which occurs after neurogenesis. Although it is not well understood how astrocytes develop from neural stem cells, it is believed that they follow a similar pattern as neurons. Astrocyte precursors are formed and migrate from stem cells. A maturing postnatal astrocyte is formed, from which an adult astrocyte develops. Local conditions and patterns help them develop heterogeneity (174, 192, 200-205).

Normal aging is known to be accompanied by mild inflammation, reactive gliosis and oxidative stress. In AD, astrocytes show some effects similar to microglia, but also have beneficial properties. They help in removing A β without the release of excessive opsonins or cytokines. After activation, astrocytes also release helpful growth factors such as nerve growth factor (NGF), S100 β (neurotrophic signaling molecule), brain derived neurotrophic factor (BDNF) and neurotrophin 3 (NT-3) (206-208). Hence they help other immune cells as well as neurons to control the release of toxic substances under inflammatory conditions. Astrogliosis is an event observed in AD brain, represented by proliferation of astrocytes in response to neuronal cell death and A β stimulation. Chronic inflammation in AD leads to an increase in chemokines like IL-6, IL-1 β , IFN- γ which further secretes other proinflammatory cytokines (193, 209, 210).

1.8 **Microglia – Macrophages of the brain**

Microglia are the tissue-resident macrophages of the brain. They are the first line of defense for the brain against any insult or threat. The four most important functions of microglia are immune surveillance, apoptotic clearance, neurogenesis and synaptic pruning. MG share a common myeloid origin with macrophages, but have a different and distinct morphology and function (194, 211, 212). These cells develop in the bone marrow and then migrate to the brain and then adapt to their residing locations. Studies in mice have revealed that microglia originate in the yolk sac in a pU1-dependent and Myb-independent manner from erythromyeloid progenitors (EMPs). During development of the zebra fish brain, the precursor cells spread within the embryo and then, 48 hrs post fertilization, invade the brain and establish a microglial population (211-216). In neurodegenerative diseases like AD, patients suffer from microglial over activation and blood brain barrier defects arising from chronic inflammation. It is believed that microglia poorly phagocytose amyloid plaques and promote AD pathology by releasing proinflammatory cytokines. MG develop a ramified and activated phenotype around A β amyloid plaques in AD and lose their intrinsic beneficial role during the disease progression. Instead they acquire a toxic phenotype which is evident from the increasing A β build up and increase in inflammation over time (141, 170, 174, 211, 217-221).




Figure 1.3 Schematic chart diagram of microglial cell in AD

Microglia are the prime immune cells in the brain. They constantly survey their surroundings for the presence of pathogens and cellular debris. It is also responsible for maintaining the homeostasis of the brain. When microglial cells encounters misfolded or aggregated proteins like Abeta or tau in AD, it triggers an innate immune response characterised by release of inflammatory chemokines and ROS, which ultimately leads to neuronal cells death.

1.9 Various functions of Microglia

There are several important functions of microglia in the context of brain development, homeostasis and neuronal connectivity. These different functions are described below.

1.9.1 Phagocytosis of debris and foreign bodies.

Phagocytosis is the most important and basic function of microglial cells. These are the cells that attack any kind of foreign bodies or debris found in healthy brain. The MG release multiple soluble cytotoxins, neurotrophins and immunomodulatory factors that help in clearing debris by phagocytosis. Chemokines released by degenerated neurons also act as a chemo-attractant for MG to detect their site of action. Studies have revealed that MG are highly motile at any stage and have filopodia-like protrusion used for continuous surveillance of the microenvironment (222-224). In a healthy brain, to facilitate successful repair of the microenvironment, MG create an environment necessary for regeneration. In order to do so, they release numerous cytokines, ROS/RNS, chemokines and other soluble mediators at the spot of repair. These chemicals attract phagocytotic and repair helper cells, which in turn contribute to tissue repair. This kind of repair activity has been observed in remyelination of damaged axons (131, 172, 174, 225, 226).

For the purpose of phagocytosis, MG express two major classes of receptors, toll like receptors (TLRs) and receptors that recognize phosphatidylserine (PS). TLRs are mainly responsible for phagocytosis of pathogens and any unnatural bodies in the brain (227-230). This process leads also to pro-inflammatory responses and the release of

tumor necrosis factor α (TNF- α), interleukin-1 (IL-1), NO and ROS. Other groups of phagocytotic receptors in MG include scavenger receptors, RAGE, Scara receptors, Fc γ receptors and integrins (54, 148, 151, 225, 231-237). These receptors are discussed in details later on.

1.9.2 Engulfment of apoptotic cells.

Apoptosis or programmed cell death is an essential step in brain development. Microglia play a significant role in removing apoptotic neurons during CNS development and in the adult brain. Studies suggest that in brain, apoptotic cells express “eat-me” signals that allow MG to recognize these cells and eventually remove them. Phosphatidylserine (238-240) (PS) is such a signal and binds PS receptors including milk fat globule epidermal growth factor 8 (MFG-E8) present on the microglial surface. The level of expression of PS determines the phagocytic behavior of MG. In zebrafish, it has been observed that brain-specific angiogenesis inhibitor 1 (BAI1) and T-cell immunoglobulin mucin receptor 4 (TIM4) proteins are required by microglial cells to perform phagocytosis of apoptotic cells. BIA1 promotes phagosomal cup formation whereas TIM4 helps in maintaining the structure of the phagosomes (241). Other signals like adenosine triphosphate (ATP) or lysophosphatidylcholine attract the microglia toward apoptotic neurons (241). In addition, apoptotic neurons release the chemokine fractalkine (CX3CL1) into the environment to send a “find-me” signal for the microglia (241). Another interesting signal that helps microglia in finding apoptotic cells are generated due to changes in Ca^{2+} . At the site of injury, these spike Ca^{2+} signals lead to the release of ATP (80, 242). The resulting ATP gradient activates the P2y₁₂ purinergic receptor (243, 244) on microglial cells, attracting them to the site of the injured neuron. MG also are

known to trigger neuronal cell death during cerebellum and hippocampus development. This process is known as “phagoptosis” and it involves the killing of Purkinje neurons by production of reactive oxygen species and superoxide anions. These ROS are produced when CD11b integrin and DAP12 are activated in MG. Studies have also revealed that release of TNF- α in activated microglia hastens the process of phagoptosis (245, 246).

1.9.3 Participation in neurogenesis and brain wiring.

MG had been considered non-functional during neurogenesis for decades. Recently, studies have revealed that they actually play an important role in neurogenesis. MG produce or support the production of neurotrophic factors such as insulin growth factor 1 (IGF-1) to promote nerve growth and survival during neurogenesis (247). Interestingly, different signaling molecules are released by microglia during development based on their location. For example in layer V, the IGF-1 signaling pathway is activated by MG whereas in the subventricular zone, interleukin 1 beta (IL-1 β) and interferon gamma (IFN γ) pathways are activated (169, 248-255). MG promote wiring of the brain, specifically at the regions close to dopaminergic and neocortical interneurons. Apart from a direct role in neurogenesis, some location specific MG also impact proliferation of neuronal precursor cells (NPC). The activation of MG inversely correlates with the number of NPC and directly impacts glial proliferation. Hence the fewer activated MG there are, the more NPC are found in the CNS (256-260).

1.9.4 Maintenance of synaptic pruning.

Synaptic pruning is one of most interesting functions of MG. Synaptic pruning is a process where synapses are remodeled during early and late postnatal stages of neuronal connections. In postnatal mouse brains, microglial synaptic pruning is observed in

hippocampus, cerebral cortex and thalamus (261, 262). The complement system, which is considered an integral part of the infection fighting process, plays an important role in brain wiring and synaptic pruning. C1q and C3 complement receptors in particular are involved in synaptic pruning through the removal of synapses through phagocytosis. It is suggested that neuronal activity is continuously sensed by microglia in the brain (263-266).

1.10 Microglial cells in Alzheimer's disease

AD is strongly associated with a state of chronic inflammation in the brain, a process in which MG play a significant role. The role of MG in AD became clearer when *TREM2* (gene encoding the TREM2 receptor) and the gene coding for its associated protein, TYROBP/DAP12, were found to be directly associated with late onset Alzheimer's disease through GWAS studies. TREM2 is a major phagocytotic receptor in MG. TREM2 is essentially required for phagocytosis of debris and A β plaques in AD. It also controls the inflammatory response of MG and helps in promoting survival and repair pathways in AD (18, 19, 226, 231, 267, 268). In addition, mutations in the microglial gene colony stimulating factor 1 receptor (CSF1R) play an important role in neurodegeneration. CSF1R is required for normal growth of resident microglia in developing brain. In EAD, CSF1R helps in microglial activation that promotes phagocytosis of A β plaques and TAU fibrils. Interestingly, in LAD CSF1R upregulation promotes over-activation of MG leading to an increase in inflammation (269, 270).

In AD, aggregation of beta amyloid plaques and tau fibrillary tangles within cells as well as in extracellular regions leads to over activation of microglia and neuronal cell death. Evidence suggests that A β fibrils attract MG. MG elevate expression of MHC class II, a sign of activation. In vivo studies in AD transgenic mice have shown that when MG come in contact with A β plaques and tau tangles they release numerous immune molecules including granulocyte-macrophage colony-stimulating factor (GM-CSF), IL-1, IL-12, IL-23 and complement proteins. In vitro studies have implicated additional cytokines and chemokines including tumor necrosis factor (TNF- α , IL-1b, IL-6, macrophage inflammatory protein (MIP)1a, macrophage chemotactic protein (MCP)-1, IL-8, macrophage-colony stimulating factor (M-CSF) and IL-12 as factors released by activated MG. Activation of the complement cascade (C1qB, C3 and C4) is also observed. Microglial specific receptors CD14, CD36, CD47, α 6 β 1, class A scavenger receptor, TLRs and receptor for advanced glycation endproducts (RAGE) are involved in the initiation of these inflammatory pathways. All of these factors in a combined manner contribute to neuroinflammation in AD (56, 91, 148, 172, 209, 225, 248, 271-276). Complexities arise when we try to define the role of such neuroinflammation in AD brain. Evidence suggests that this inflammatory response is beneficial to the brain in the early stages of disease progression, but prolonged chronic inflammation forces MG to attain an altered state that causes more harm than good. MG at this stage continue secreting neurotoxic chemokines, cytokines, ROS, RNS and superoxide ions which worsens the AD pathology by increasing stress in the microenvironment (75, 141, 177, 178, 182, 217, 277-281).

Like other cells of myeloid origin, MG perform phagocytosis with high efficiency in healthy brain. In AD brain, MG are supposed to phagocytose A β and degrade A β by microglial proteases, which are key mechanisms for removal of these aggregates from the brain. Studies have indicated that in moderate or later severe stages of AD, the phagocytosis of A β plaques and fibrils by MG is impaired. The increasing burden of A β plaques in the AD brain increase chronic inflammation and leads to cellular senescence (219, 282, 283). Senescence in MG results in a state where MG cannot phagocytose but generates pro inflammatory factors, cytokines and reactive species. Soluble A β (sA β) has been reported to be engulfed by MG through both phagocytosis and pinocytosis. Pinocytosis is an endocytic process where MG uptake soluble A β by forming an invagination. The A β is then suspended in small hollow vesicles. The content of these vesicles are further hydrolyzed in lysosomes and degraded (284, 285).

Genetic mutations in microglia have been found to promote AD pathogenesis. For example, the TREM2 adaptor molecule, TYROB/DAP12 is associated in LOAD pathogenesis. TREM2 is one of the major receptors required by MG for phagocytosis of debris and foreign bodies. DNAX-Activation Protein 12 (DAP12) is an adaptor receptor protein that helps in ITAM receptor signaling in MG. It also interacts with CSF1R in Hereditary Diffuse Leukoencephalopathy with Spheroids (HDLS) pathology (16, 21, 231, 286, 287). Genome wide association studies (GWAS) studies revealed an increase in microglia of CD33 (Siglec-3) in human AD brain, which is correlated with increased insoluble A β levels. CD33 receptors bear an ITIM that is required for receptor signaling in MG. Increase in CD33 activity in AD mouse models revealed reduced uptake A β clearance by MG. This suggests that CD33 activity inhibits phagocytosis ability of MG

through ITIM receptor signaling (14, 288-290). TREM2 on the other hand is involved in cytokine and chemokine production, phagocytosis and MG cell motility. These studies on microglial specific genes have provided us with deeper insight in understanding the role of MG in AD progression and specifically, sporadic AD.

1.11 **SYK kinase and its activation**

SYK is a non-receptor protein-tyrosine kinase located in the cytoplasm. SYK was initially discovered as a 40 kDa catalytic fragment by two independent groups as published in Zioncheck et al. and Kobayashi et. al. Later, the full-length 72 KDa enzyme was identified through the use of specific antibodies (291, 292). SYK is mainly expressed in spleen and thymus and is found in many hematopoietic cells including B cells, thymocytes, erythrocytes, macrophages and platelets (271, 293, 294). SYK also has been found in a subset of non-hematopoietic cells including epithelial, endothelial, fibroblast, neuronal cells and some non-neuronal brain cells (295, 296). SYK is well known to play crucial roles in adaptive immune receptor signaling, innate immune signaling, platelet activation, cellular adhesion, cellular motility, vascular development, epithelial cell division, and osteoclast maturation (297, 298). SYK function has been linked to several disorders including autoimmune diseases, hematological malignancies, rheumatoid arthritis and inflammation (299-306).





Figure 1.4 Schematic diagram of SYK structure

SYK is a protein tyrosine kinase which consists of a C-terminus catalytic domain to promote activation of downstream proteins. It also has two tandem SH2 domains at the N-terminal that helps in receptor signaling through ITAM binding.

As a kinase, SYK phosphorylates multiple proteins on tyrosine to control key receptor signaling pathways including Ca^{2+} mobilization and the phosphoinositide 3-kinase (PI3K)/AKT, NF- κ B and RAS/ extracellular-signal-regulated kinases (ERK) pathways (307-313). At least two substrates of Syk, CD79a and the Ikaros protein, also have been reported to be phosphorylated by SYK on serine, suggesting that SYK may have capabilities beyond catalyzing tyrosine phosphorylation. Its ability to couple with immune recognition receptors makes it an important kinase in the immune system that regulates multiple pathways in response to many extracellular signals including antigens and immunoglobulin-antigen complexes (314).

The N-terminus of SYK consists of two 100 amino acid conserved sequences known as Src Homology 2 (SH2) domains. These SH2 domains are connected by a region known as linker A and both are then separated by linker B from the C-terminal kinase domain. These domains bind to proteins containing phosphotyrosine and thus initiate protein-protein interactions. The two SH2 domains of SYK bind specifically to proteins that contain immunoreceptor tyrosine-based activation motifs (ITAMs). ITAMs are characterized by two short peptide sequences of YXXL/I (where Y is tyrosine, L and I are leucine and isoleucine, respectively, and X denotes any amino acid) separated by a spacer of 6-12 amino acids. Once receptors with ITAMs are oligomerized upon engagement, the two tyrosines are rapidly phosphorylated leading to the recruitment and activation of SYK (23, 291, 304, 315-317). In some cells, particularly T cells, the SYK-family kinase ζ -chain-associated protein kinase of 70 kDa (ZAP70) is recruited to phosphorylated ITAMs (318, 319). Following its activation, multiple tyrosines on SYK become phosphorylated including sites within the linker A and B regions along with

tyrosines within the catalytic domain and at the extreme C-terminus. SYK both autophosphorylates and is phosphorylated in *trans* by LYN. Examples of important phosphorylation sites include Y130 located in linker A, Y317, Y342 and Y346 located in linker B, Y519 and Y520 located on the activation loop in the catalytic domain and Y624 and Y625 near the C-terminus. Phosphorylations serve to regulate the activity of SYK, to modulate kinase-receptor interactions and to promote protein-protein interactions with downstream effectors containing SH2 or related domains (304, 320, 321).

Following its activation, a fraction of SYK is released from the receptor and can shuttle between the nucleus and the cytoplasm to promote further downstream signaling events. Included among the SYK substrates that have been well characterized are the B cell linker protein (BLNK/SLP65), SH2 domain leukocyte-specific phosphoprotein 75 kDa (SLP76), linker for activation of T cells (LAT), PI3K, phospholipase C γ (PLC γ), Bruton's tyrosine kinase (BTK), Vav-1 (Ras/Rho guanine nucleotide exchange factor), β -tubulin, and CBL (E3 ubiquitin ligase) (322-330).

1.12. **Receptor based immune signaling of SYK**

1.12.1 **Syk in B cell signaling.**

A major component of the adaptive immune response in the body requires B cell activation for producing antigen specific antibodies to target foreign molecules and pathogens. It is the B cell receptor for antigen (BCR) that recognizes foreign antigens. The BCR consists of a membrane-bound immunoglobulin associated with a heterodimer of CD79a and CD79b subunits. CD79a and CD79b also are commonly known as Ig- α

and Ig- β . The BCR contains ITAM sequences in the cytoplasmic tails of CD79a and CD79b that are initially phosphorylated by a member of the Src-family of protein-tyrosine kinases. The main Src-family kinase in B cells is LYN, which phosphorylates the first tyrosine of the ITAM. The second tyrosine is speculated to be phosphorylated by SYK based on in vitro studies. Once both of the ITAM tyrosines are phosphorylated, SYK binds tightly through its tandem SH2 domains. Once bound to the BCR, active Syk phosphorylates additional ITAM tyrosines leading to additional SYK binding to the BCR complex. Activated SYK then catalyzes the phosphorylation of the multiple protein substrates that play important roles in antigen receptor signaling to regulate downstream cell signaling pathways. These include for example, the adapter protein BLNK, whose phosphorylation creates docking sites for BTK and PLC- γ to form a complex ultimately facilitating calcium mobilization (315, 331-336).

1.12.2 Syk in mast cell signaling.

Mast cells and basophils are key players in inflammation. Their activation results in type I allergic reactions through the release of pre-formed inflammatory mediators like histamine from cytoplasmic granules and the generation of additional biologically active mediators like leukotrienes. Mast cells are enriched with high affinity IgE receptors (Fc ϵ R1) on their cell surface (23, 291, 317, 337-340). The stimulation of mast cells is initiated by the binding of antigens to Fc ϵ R1 receptors pre-associated with IgE. Fc ϵ R1 is a tetrameric protein consisting of $\alpha\beta\gamma_2$ chains. The Fc region of IgE binds to the α -chain of IgE. ITAMs are present in the β - and γ - chain, located in their cytoplasmic domains. Similar to the BCR, Fc ϵ R1 does not have an intrinsic enzymatic activity, but relies on the activation of non-receptor protein-tyrosine kinases for its activity (311, 341-343).

LYN initiates the phosphorylation of the tyrosines on these ITAMs leading to the docking of SYK and its activation. Active SYK further propagates downstream tyrosine phosphorylation of adaptor proteins such as LAT1, LAT2 and SLP76. Several other substrates including PI3K, BTK, PLC γ 1, PLC γ 2 and PKC are subsequently activated through this pathway. Studies indicate that the absence of SYK in mast cells leads to a complete loss of IgE-mediated degranulation and release of cytokines, making SYK a key mediator of mast cell signaling and a popular drug target (297, 305, 317, 344, 345).

1.12.2 Syk in macrophage signaling.

Macrophages are crucial components of the innate immune response and are the first line of defense against pathogens. The most important function of macrophages is to recognize foreign substances to either engulf and degrade them or to engage other immune cells to destroy them. They are white blood cells of the myeloid lineage and differentiate in tissues from circulating monocytes. During inflammation, macrophages perform vital functions including phagocytosis, antigen presentation and immunomodulation (271, 346-349). The activation of macrophages occurs through various stimuli including cell surface components of pathogens such as lipopolysaccharide (LPS), cytokines such as interferon- γ and TNF- α , apoptotic cells and TH1 T cells. Macrophages express a broad range of receptors including the toll-like and dectin-1 pattern recognition receptors, scavenger receptors, and Fc γ receptors that recognize foreign substances and opsonized particles in the body (125, 350, 351). Many of these receptors are coupled to downstream signaling pathways through protein kinases that, in turn, activate transcription factors such as nuclear factor- κ B (NF- κ B), activator protein-1 (AP-1), and cAMP response element binding protein (CREB) that promote the

expression of proinflammatory genes. Many of these receptors also trigger phagocytosis, the secretion of inflammatory mediators, the generation of ROS, the production of nitric oxide and the generation of arachidonic acid metabolites such as prostaglandin E₂ (PGE₂) (137, 337, 352-361).

Interestingly, many of these receptors in macrophages contain ITAM sequences and are thus dependent on SYK for their activities. These include the Fc γ receptors, which are mainly involved in the phagocytosis of foreign bodies to which IgG is associated. These belong to the immunoglobulin gene superfamily. Fc γ receptors are classified into three categories: class I and III, which form multimeric complexes, and class II receptors that are monomeric. All Fc γ receptors contain an extracellular binding domain that recognizes the Fc region of IgG. Macrophages primarily express activating Fc γ receptors that stimulate phagocytosis and other innate immune responses. Clustering of these receptors leads to the Src-family kinase-mediated phosphorylation of tyrosines within ITAMs found either within the receptor or receptor complex. Subsequently SYK is recruited to the phosphorylated ITAMs and activated, resulting in the phosphorylation of other Fc γ receptor ITAMs and downstream substrates. Fc γ RI, Fc γ RIIA, and Fc γ RIIIA receptors specifically bind to SYK and their clustering leads to its activation. Several studies in primary macrophages have revealed that the absence of SYK leads to a failure of Fc γ receptor-mediated phagocytosis and a loss of the receptor stimulated generation of NO and ROS, events that are normally enhanced by Fc γ receptor engagement in macrophages (25, 26, 292, 315, 362-364).

LPS is a commonly used stimulus for macrophages. LPS is found on the surfaces of Gram-negative bacteria. Thus, exposure to LPS mimics a bacterial infection. The

TLR4 receptor is needed for macrophages to respond to LPS. Interestingly, SYK also is activated in macrophages by LPS stimulation. There is evidence that SYK is constitutively docked to TLR4 receptors despite the lack of an ITAM. Active Syk thereby promotes activation of downstream signaling molecules, such as AKT, PI3K, IKK, NF- κ B and phosphoinositide-dependent kinase-1 (PDK1), ultimately leading to induction of TNF- α , COX-2 and inducible nitric oxide synthase (iNOS), and the secretion of nitric oxide (NO), ROS and PGE₂. Studies also indicate that minimally oxidized low-density lipoprotein (mmLDL) activates SYK in macrophages through TLR4. Activated SYK subsequently activates Vav1, PLC γ and JNK resulting in cytoskeleton modifications, ROS generation and cytokine secretion (293, 312, 365-369).

1.12.4 Syk in neutrophil signaling.

Neutrophils are also essential cells in the innate immune system. They are the most abundant type of white blood cell and also the first cell type to reach a site of inflammation. Once in contact with cellular and humoral factors, they become primed and travel through blood vessels to the site of microbial infection. Their main function is to kill or suppress microorganisms during the innate immune response.

Integrin receptors are particularly important receptors in neutrophils and are required for cell adhesion, motility, spreading, the respiratory or oxidative burst, release of cytokines and release of antimicrobial granule proteins. β_1 , β_2 , and β_3 integrins in neutrophils all associate with SYK to mediate their functions. Neutrophils mainly express the β_2 integrins $\alpha_L\beta_2$, $\alpha_M\beta_2$ and $\alpha_4\beta_1$. Integrin signaling occurs in two steps, known as inside-out (integrin activation) and outside-in (aggregation of integrins) signaling. SYK is essentially involved on both these signaling pathways. Studies in SYK negative

neutrophils have revealed that, in the absence of SYK, neutrophils fail to spread when plated on integrin ligands and fail to undergo respiratory bursts or degranulation in response to integrin clustering. Integrins themselves lack ITAM motifs, but in neutrophils signal instead through the adaptor proteins DAP12 and FCγR, both of which contain ITAMs that, when phosphorylated, bind and activate SYK (246, 286, 303, 327, 370-372).




Figure 1.5 Schematic diagram of adaptor mediated SYK-ITAM signaling

SYK is activated through binding of receptor ITAM to SH2 domains. In receptors like BCR, it requires LYN kinase to initially phosphorylate ITAM on receptor adaptor molecule. Once activated the receptor further activates through autophosphorylation and initiation of SYK-ITAM cascade. This activation leads to activation of several downstream signaling pathways through tyrosine phosphorylation.




Figure 1.6 Schematic diagram of direct ITAM-SYK signaling

In some receptors like FcγRIIA, SYK is activated through binding of receptor ITAM present at the cytoplasmic tail of the receptor. It does not require an adaptor protein for activation. LYN kinase initially phosphorylates ITAM on receptor tail and SYK binds to phosphorylated ITAM through SH2 domains. Thus SYK-ITAM signaling cascade is initiated.

1.13 Syk in the brain

Brain cells can be classified into neuronal and non-neuronal cells. Non-neuronal cells in the brain include glial cells and other helper cells that assist with the normal functioning of the brain. Glial cells mainly consist of astrocytes, oligodendrocytes and microglia. One crucial function of glial cells is to comprise the immune system of the brain, which functions quite differently from the rest of the body. The presence of the blood brain barrier restricts the ability of normal immune cells in the blood to infiltrate into a healthy brain to maintain immunity. While the brain had long been considered a completely immunologically privileged site, interesting new data indicate the presence of a lymphatic system that allows T cells access to the meninges (259, 287, 311, 369, 373-375).

Although the role of SYK in the brain is poorly understood, considering its vital role in the majority of immune cells; SYK's role in the brain's immune system cannot be neglected.

1.13.1 Syk in neurons

Few in vitro studies have suggested the presence of SYK in neurons. It has been mentioned that SYK is involved in neuronal differentiation in P19 cells (376). Another study on the CD3 ζ adaptor immune protein in the CNS has suggested that SYK is involved in neuronal morphogenesis (377). The presence of an ITAM on CD3 ζ allows it to bind to the SYK SH2 domains thereby activating SYK. Without SYK, CD3 ζ could not promote neuronal morphogenesis. Studies on sensory neurons have reported that SYK is required for NF- κ B activation through the cytokine ciliary neurotrophic factor (CNTF).

Upon CNTF stimulation, SYK is activated and phosphorylates I κ B leading to NF- κ B activation. This pathway also promotes neurite growth from developing neurons (313, 378).

1.13.2 Syk in microglia

Microglia are the most vital immune cells in the brain. In the CNS, they are mainly responsible for phagocytosis of debris, foreign particles, apoptotic cells, and microbes. Apart from these activities, they also maintain homeostasis in the brain through the secretion of anti-inflammatory and pro-inflammatory factors. Phagocytosis by MG is mainly mediated through various receptor types including TLRs, PTRs, PAMPs, RAGE, SCARA1, SCARA2, CD36, CD68, integrins and Fc γ receptors. Many of these immune receptors contain ITAMs as discussed above in macrophages (141, 148, 225, 228, 231, 272, 379-381).

ITAM-mediated receptor signaling through SYK is crucial for phagocytosis. DRAPER, for example, is a phagocytotic receptor in *Drosophila* that binds to Shark (fruit fly version of SYK) and is important for phagocytosis. Both phosphorylation of DRAPER ITAM tyrosines and SHARK activation are required for proper phagocytosis of damaged axons and debris (259, 382). The homologs of DRAPER in mammals are JEDI1 and multiple epidermal growth factor-like domains 10 (MEGF10). In the peripheral nervous system (PNS), 50% of the neurons are eliminated through apoptosis during development. JEDI1 and MEGF10 are essential for glial cell precursors to mediate phagocytosis of apoptotic neurons during development. Both contain two ITAMs in their intracellular domains that, when phosphorylated, bind to the SH2 domains of

SYK. In the absence of SYK, engulfment of apoptotic neurons is inhibited, but phagocytic activity can be recovered when cells are reconstituted with SYK (259).

DAP12/TYROBP is an adapter protein that contains an ITAM and is important for phagocytosis by mammalian MG. DAP12 associates with both TREM2 and signal regulatory protein- β 1 (SIRP β 1). When phosphorylated following receptor engagement, DAP12 associates with SYK to mediate phagocytosis of damaged neurons, demyelinated axons and other cellular debris. This involves the binding of the SH2 domains of SYK to the phosphorylated ITAM of DAP12. TREM2 is important for the uptake of apoptotic neuronal cells. Additional studies suggest that TREM2 also binds to lipo-oligosaccharides of both Gram-positive and -negative bacteria. Binding stimulates TREM2 and, via ITAM and SYK-dependent signaling, promotes phagocytosis of bacteria. Interestingly, certain heterozygous mutations in TREM2 are correlated with enhanced incidence of AD. Furthermore, DAP12 was identified recently as a key regulator of genes associated with late-onset AD (18, 21, 231, 254, 287, 383, 384).

Immunoreceptor tyrosine based inhibition motifs (ITIMs) resemble one-half of an ITAM. Receptors bearing ITIMs negatively regulate phagocytosis. ITIMs inhibit ITAM-mediated signaling by recruiting the SH2 domain-bearing protein tyrosine phosphate SHP1 and the lipid phosphatase SHIP1 to counteract the activation of protein and lipid kinases that occurs during ITAM-mediated signaling (21, 385). CD33 related sialic acid-binding immunoglobulin superfamily lectins (SIGLECS) are examples of proteins bearing ITIMs. They belong to the Ig superfamily and recognize specific sugar residues on cell surface glycans. Since they contain an ITIM sequence in their cytoplasmic domains, they negatively regulate ITAM-SYK receptor signaling in MG. Interestingly,

CD33 expression is enhanced in AD brains and has been linked genetically to AD in GWAS studies (20, 288, 289, 386).

Another group of receptors that play a vital role in the phagocytotic ability of MG are Fc γ receptors. As discussed above, many of these contain ITAMs and thus require ITAM-SYK mediated signaling for engulfment of opsonized particles. Clustering of Fc γ receptors initiates ITAM phosphorylation, which in turn docks SYK through its SH2 domains. This ITAM-SYK interaction further activates SYK and hence activates downstream signaling pathways to facilitate phagocytosis of opsonized particles and antigens (9, 273, 387-389).

DECTIN-1, a major beta-glucan receptor that promotes phagocytosis of fungal pathogens, also is associated with SYK in MG. DECTIN-1 is unusual in that it's cytoplasmic tail contains only one-half of an ITAM (known as a hemITAM). DECTIN-1 is thought to function as a dimer whereby two DECTIN-1 molecules bind simultaneously to a single SYK molecule to promote its activation (223).

Several studies have revealed that SYK is essential not only for phagocytosis by MG, but also for cytokine production, NO synthesis and ROS generation. SYK mediates IL-1 β and TNF- α secretion in MG through activation of NF- κ B. The secretion of these cytokines into culture media can promote apoptosis of co-cultured adult neurons. β -amyloid fibrils or prion protein aggregates can activate SYK in MG cells. This activation is accompanied by intracellular calcium release and PKC activation leading to the production of IL-1 β and TNF- α (210, 221).

1.14. **Stress granules and their components**

SGs are cytoplasmic RNA-protein complexes that appear in eukaryotic cells in response to various kinds of stress-related stimuli that impair translation. Some of these stressors include heat, viral infection, oxidative stress, ultraviolet irradiation, and hypoxia. SGs were first identified in cells cultured at high temperature, which led to the formation of cytoplasmic granules containing heat shock proteins. The phosphorylation of eukaryotic translation initiation factor 2- α (eIF2 α) is one of the initial steps in SG assembly induced by stress. Based on the stress stimulus, distinct eIF2 α kinases can be activated ((protein kinase R (PKR), protein kinase RNA-like endoplasmic reticulum kinase (PERK), Heme-regulated eIF2 α kinase (HRI) and general control nonderepressible 2(GCN2)) (390-392). The subsequent phosphorylation of eIF2 α stalls assembly of ribosomes at the 48S pre-initiation complex. As a result, this initiation complex remains assembled at the 5' end of mRNAs resulting in translation impairment and a block in polysome assembly. These nonpolysomal transcripts are then aggregated into complexes along with many other RNA binding proteins to form SGs (393, 394).

Since their discovery, numerous studies have shown that SGs contain many biological components important for mRNA metabolism. The major components include nontranslating mRNAs, translation initiation complexes, RNA binding proteins, mRNA processing helper proteins, aggregated proteins, kinases and other proteins involved in stress signaling. Some of the major proteins involved in SG formation are: AU rich element binding proteins, Argonaute, eIF2 α , fragile X mental retardation protein (FMRP), fragile X mental retardation related protein 1 (FXR1), Ras-GTPase-activating protein

SH3-domain-binding protein (G3BP), glycine-tryptophan 182 (GW182), poly(A)-binding protein 1 (PABP1), T cell internal antigen-1 (TIA-1) and T cell internal antigen-1-related (TIAR) (395-398).

Recent studies have shown that many proteins responsible for RNA splicing, transcription, adhesion, immune signaling, autophagy and apoptosis also associate with SGs. The function of SG formation in various biological systems or conditions is yet to be completely deciphered. Studies suggest sequestration of mRNAs in SGs is a way to protect them from stress and conserve energy to overcome stress. SG are highly dynamic in nature and self-assemble and disassemble based on changes in the environment of the cell (28, 399).

1.14.1 SG assembly and disassembly

The presence of diverse proteins, RNAs and other biological molecules suggest that SG assembly is very dynamic and variable depending on the biological system on the type of stress induced. In some instances, composition of SGs depends on the type of mRNPs present in the SG at that specific stress stimulation. Since specific isolation of SGs has not been possible yet, most of the studies on SG are done through microscopy or correlated to the function of SG components under stress conditions. Such studies have reported that the size and shape of SG can vary in different systems. It also mentions that SGs do not confer to a definite structural organization (398, 400-403).

SG assembly is mainly modulated by three factors: covalent protein modifications, protein-protein interactions and microtubule network dynamics. In terms of protein modifications, a key process is phosphorylation. As described above, eIF2 α phosphorylation is one of the most crucial steps in SG assembly. Other proteins that

undergo phosphorylation during SG assembly are TRISTETRAPROLIN (TTP), B-related factor 1 (BRF1) and Ras-GTPase-activating protein SH3-domain-binding protein (G3BP). In addition, the acetylation of histone deacetylase 6 (HDAC6) also has been reported to affect SG assembly as has protein methylation and methyl group binding to proteins with Tudor domains. One study also mentioned the role of ubiquitin-dependent proteasome system (UPS) in SG assembly. UPS interference modulates mRNA decay. It promotes phosphorylation of eIF2 α and sequestration of AU-binding proteins (AUB) and ARE in SGs thereby preventing their decay under stress (15, 27, 398, 404).

Recent studies have shown that protein-protein interactions are critical for SG assembly. Many RNA-binding proteins like G3BP contain dimerization domains that promote protein-protein interactions that aid in SG formation. In fact, the overexpression of G3BP alone is sufficient to induce SGs. Additionally, RNA metabolism helper proteins like TIA-1 that contain QN-rich prion-like domains also participate in SG assembly through self-aggregation. Examples include the prion-like proteins TIA-1 and TIAR, which accumulate in the cytoplasm as complexes under SA or MG132 mediated stress promoting SG assembly. Proteins susceptible to unfolding or misfolding ((TAR DNA-binding protein 43 (TDP-43), Fused in Sarcoma (FUS) and TAU)) observed in several diseases like ALS and AD also bind to each other along with other RNPs to promote SG assembly. A third important factor that contributes to SG assembly is the microtubule network. Microtubules associated motor proteins like dynein and kinesin also associate with SGs and regulate their assembly and disassembly (398, 405-407). SGs disassemble following removal of the stress. Stress removal repairs the translation initiation machinery and helps SG components to disassociate to allow mRNAs to resume

translation or to be transferred to processing bodies for degradation. Heat shock proteins that disaggregate prion like-proteins have been correlated with SG disassembly. The RNA-binding protein Staufen1 (STAU1) has been associated with SG dynamics. The overexpression of STAU1 promotes SG disassembly whereas knockdown promotes assembly. Studies suggest that STAU1 promotes the stabilization of the mRNA-polysome complex during this process (408). Another study showed that, upon stress removal, FAK promotes the phosphorylation of growth factor receptor-bound protein 7 (GRB7) to weaken its interactions with SG component proteins like TIA-1, leading to SG disassembly (409).

Autophagy has been recently linked to SG removal. It has been shown that SG clearance is reduced in eukaryotic cells by inhibition of autophagy or depletion of proteins like valosin-containing protein (VCP) and autophagy related protein (ATG) which is required for autophagy machinery in cells (32).

1.14.2 SG components

Depending on the cell type, stress induced and biological system under study, the components of SG can vary. The first class of SG components represents members of translation initiation complexes. Major proteins in this category include eIF2 α , eIF3, eIF4F, eIF4B, components of small ribosomal units and PABP-1. These are considered core components of SGs. The second class of components are the mRNA binding proteins, translation modifying proteins and proteins involved in mRNA stability. Few examples of such proteins are TIA-1, TIAR, FMRP, FXR1, FAST, ARGONAUTE, cytoplasmic polyadenylation element binding protein (CPEB), ATAXIN-2, and RNA-associated protein 55 (RAP55). Proteins involved in RNA decay that are SG components

include TTP, BRF1, receptor for Activated C Kinase 1 (RACK1), Z-DNA Binding Protein 1 (ZBP1) and PMR1 (27, 28, 394, 406, 410).

Finally, the third class of SG components consists of RNA binding protein that controls RNA decay and processing. In this category we have G3BP, CAPRIN, FAST, survival of motor neuron (SMN) and Ins (1,3,4,5,6)P5 2-kinase (IP5K). Few of these SG components like TIA-1, G3BP can be a part of both second and third class. Recent studies have shown that some proteins that do not directly interact with RNA, interact instead with core SG components to become a part of a SG. For example, steroid receptor coactivator-3 (SRC-3), TAU, folate-binding protein (FBP) proteins are found in SGs because of their ability to interact with TIA-1; TNF Receptor-Associated Factor 2 (TRAF2) on the other hand binds to eIF4G and DIS1 binds to eIF3 (29, 411-413).

1.14.3 SG functions

The true function of SGs in the cell is not totally clear. Studies indicate that SGs are present in very different biological systems and participate in multiple processes. SG components have roles in biological functions like mRNA stability, mRNA translation and mRNA modification outside of their localization in SGs. Hence knockdown studies of SG components do not directly address the function of SGs. One of the major suggested consequences of SG formation is to regulate mRNA stability. Formation of SGs containing untranslated mRNAs under stress protects them from degradation in the cytoplasm. Once the stress is gone and SG disassembly occurs, these mRNAs are freed to participate normally in translation. Since SGs consist of mRNP complexes including translation initiation complexes the formation of SGs under stress may assemble and store specific mRNP proteins for participation in future translation processes. Since under

stress translation is stalled, excess mRNPs required for translation are stored as SGs which are later utilized once the stress is gone (398, 404, 414-416).

Under stress, a cell ultimately needs to decide if it will be able to recover from the stress or, if the stress is too extreme, to undergo apoptosis through programmed cell death. SGs play important roles in the apoptosis of cells under stress. For example, MAP kinase kinase kinase (MTK1) is activated in cells under severe stress. MTK1 activation-mediated apoptosis takes place through an interaction with receptor for activated C kinase 1 (RACK1). In vitro studies have shown that, under stress, RACK1 is sequestered to SGs, thereby inhibiting MTK1 activation and preventing apoptosis (417, 418).




Figure 1.7 Schematic diagram of stress granules in neurodegeneration

In neurodegenerative diseases like AD, ALS and SMA, presence of SGs have been reported in human brain and animals models. Core SG components like G3BP1, TIA-1 associate with misfolded and aggregated proteins like TAU, TDP-43 in neurons. The size and number of stress granules increases with the disease progression. Possible role of stress granules in neurodegeneration is yet unknown.

1.15. Stress granules in different neurological diseases

1.15.1 SGs in Alzheimer's disease

AD is a progressive neurological disorder. As mentioned earlier, oxidative stress is one of the key players in AD. Protein misfolding and accumulation of tau proteins and A β is a common event in AD. As a consequence, chronic neuroinflammation and neuronal cell death is observed. Studies have suggested that SG components like TIA-1, TTP and G3BP associate with pathogenic tau epitopes. TIA-1 positively regulates apoptotic cell death and regulates mRNA translation of TNF- α and COX-2. TTP on the other hand associates with mRNA for TNF- α , GM-CSF, COX-2, IL-3, IL-10, and interferon- γ making it an important protein in inflammatory pathways (29, 419, 420).

1.15.2 SGs in Amyotrophic Lateral Sclerosis (ALS)

ALS is a neurodegenerative disease that mainly affects the motor neurons. It is considered the most common motor neuron disease. The neuronal death mainly takes place at the brain stem, spinal cord and motor cortex. As a result of such degeneration, muscle fasciculation, wasting and weakness; increased spasticity, and hyperreflexia are observed. One of the hallmark proteins involved in ALS is TDP-43. In ALS patients, TDP-43 is mutated, mis-folded and aggregated in degenerating motor neurons. Recent studies have shown that TDP-43 is localized in SGs in ALS. Another disease that is closely linked with ALS is Frontotemporal Lobar Degeneration (FTLD). TDP-43 is also mutated in FTLD. TDP-43 is mainly a nuclear protein, but in neurodegenerative diseases it is mis-localized to the cytoplasm. TDP-43 plays an important role in mRNA metabolism. In neurons, it mediates splicing and mRNA stability. Hence it influences proteins that are

required for proper formation of synapses in neurons. Since TDP-43 is localized to cytoplasmic SGs in ALS, it cannot further maintain mRNA stability. On the other hand, TDP-43 has a self aggregating domain and an excess of TDP-43 promotes aggregation. Such aggregation of TDP-43 in turn elevates its expression. Excess TDP-43 has been reported as neurotoxic to neurons and promotes apoptosis. TDP-43 tightly binds to G3BP in neurons. This binding promotes self aggregation of TDP-43 leading to larger SGs. Other proteins that localize to SGs in ALS are FUS, HUNTINGTIN and PrP (38, 405, 406, 421-423).

1.15.3 SG in spinal muscular atrophy

SMA is a recessive neuromuscular disorder that affects the α -motor neurons. It mainly occurs in infants and involves the spinal cord. Mutations in or deletions of the telomeric copy of the *SMN-1* gene and retention of the telomeric copy of *SMN-2* characterizes this disease. SMN plays a key role in axonal growth and maintenance through regulation of mRNA metabolism. SMN has been reported to be localized in SGs. SMN localization in SGs impairs its ability to regulate protein translation leading to neuronal cell death (412, 424, 425).

CHAPTER TWO: MATERIAL AND METHODS

2.1 Cells and cell lines

BV-2 microglial cells developed by Blasi and colleagues (387) were obtained from Dr. Chris Rochet (Purdue University) and cultured in high glucose DMEM media containing 10% FBS and 100 U/ml penicillin and streptomycin. The cells were passaged at a ratio of 1:8 every 48 h and a confluency of 60-80% was maintained at all times. Fresh cells were passaged at least two times before conducting an experiment. The use of freshly thawed cells was limited to 12 passages.

N9 microglial cells⁽⁴²⁶⁾ were obtained from Dr. Fabio Bianco (Neuro-Zone, Italy) and were cultured in ISCOVE medium containing 10% FBS and 100 U/ml penicillin and streptomycin. Cells were passaged at a ratio of 1:6 every other day to obtain consistent results. The confluency of these cells was maintained at 50-70 % at all times. Freshly thawed cells were used for a maximum of 15 passages.

HEK293T cells were purchased from ATCC and HT22 mouse hippocampal neuronal cells⁵⁸ were obtained from Dr. Shaohua Yang, (University of North Texas Health Sciences Center). Both were grown in DMEM containing 10% FBS and 100 U/ml penicillin and 100 mg/ml streptomycin. All cell lines were grown at 37°C in 5% CO₂.

2.2 Plasmids and Lentiviral vectors

The cDNA coding for mouse Syk-EGFP fusion protein was cloned into a puromycin resistance lentiviral vector pLVX as described (427). Lentivirus particles were generated by cotransfecting HEK293T cells (2×10^6 cells/10 cm plate) with 5 μ g of Syk-EGFP pLVX, 5 μ g of pHR' CMV-R8.20 VPR and 2.25 μ g of pHR' CMV-VSVG using Lipofectamine 200 (Invitrogen). After 48 h the media supernatant containing viral particles was collected and filtered using 40 μ m pore size filter. Harvested lentiviral particles were added to BV-2 cells (2×10^5 cells/10 cm plate) in the presence of 10 μ g/ml polybrene. 48 h post infection, cells were selected in the presence of 5 μ g/ml puromycin for 92 h. Finally, cells were maintained in 2.5 μ g/ml puromycin and passaged every 48 h.

To establish the mouse Syk knockdown BV-2 cell line, BV-2 cells (2×10^5 cells/10 cm plate) were infected with one of a set of six independent GIPZ mouse Syk shRNA lentiviruses (GE Dharmacon). A scrambled shRNA was also infected to establish a control cell line. Six independent Syk shRNA and scrambled shRNA lentiviral particles were generated as described above with the lentiviral vectors for Syk-EGFP.

Lipofectamine 3000 (Invitrogen) was used instead of Lipofectamine 2000 to transfect HEK293T cells. One of the plasmids among the six exhibited robust knockdown as evaluated by western blotting. Syk shRNA #5 plasmid was used to conduct the future experiments.

2.3 **Primary cell culture and animal models**

Primary microglial cells were isolated essentially as described (428, 429) from 1 month or 20 month old wild-type or *Syk*^{+/-} C57BL/6 mice. *Syk*-haplosufficient mice were generated by gene targeting by Ingenious Targeting Laboratory, Ronkonkoma, NY. Approximately, 5 mice from each group were anesthetized using isoflurane anaesthetizing equipment and perfused in the left ventricle at constant pressure for 5 min with ice cold Ringer's solution containing 2 U/ml heparin (Cat. No. AK3004, Akron Biotech, USA). Immediately brains were aseptically removed and meninges separated. The brain was minced and dissociated with a sterile scalpel blade in media containing 1 mg/ml papain, 1.2 U/ml dispase II (Cat. No. P4762 and D4693, Sigma-Aldrich, USA) and 20 U/ml DNase1 (Cat. No AK3778, Akron Biotech, USA). Following pH neutralization of media, cells were pelleted and recovered, pipetted up and down using glass Pasteur pipettes of decreasing hole sizes, filtered through 70 μ m and 40 μ m cell strainers and collected in media. Cells were separated on a discontinuous percoll (Cat. No. 17-0891-01, GE Healthcare, USA) density gradient, collected, washed, and cultured on 1 μ g/cm² fibronectin (Cat. 341631, Calbiochem)-coated dishes. Three different concentrations (70% SIP, 37% SIP and 30% SIP) of saturated isotonic percoll (SIP) were used and microglia were present between 37% and 70% SIP. Isolated cells were washed and cultured in DMEM containing 5 ng/ml murine carrier free granulocyte and macrophage colony stimulating factor (GM-CSF) (Cat. No. 130-094-043, Miltenyl Biotech), 10% FBS and 100 U/ml penicillin and 100 mg/ml streptomycin. For experiments, 1 X10³ cells/well in a 24 well plate were grown in high glucose DMEM

containing 10% FBS and 100 U/ml penicillin and 100 mg/ml streptomycin. Primary microglia were confirmed by western blotting and immunofluorescence staining of IBA1 (428). All mice were housed and experiments were done in accordance to the guidelines of IACUC at Purdue University.

2.4 **Stress granule formation**

BV-2 or N9 cells (1×10^4 cells/well in a 24 well plate) were grown on poly-D-lysine coated cover slips in media containing 1% FBS for 12 h before treatment with sodium arsenite (SA) (1 μ M), A β (1-42) (100 nM), or A β (1-42) fibrils (100 nM) for the indicated times. A β (1-42) (rPeptide, Cat. No. A-1008) or A β (42-1) (American Peptide Company, Cat. No. 62-0-81) were resuspended in DMSO, 1% NH₄OH at a concentration of 0.025 M. To prepare A β fibrils, DMSO soluble A β was sonicated for 30 min at 1 min intervals, shaken at 37°C for 48 h and then at room temperature for 72 h (59, 430). Treated cells were fixed in 4% formaldehyde in PBS, permeabilized in 1% Triton X-100, blocked with 5% goat or donkey serum (depending on the species of origin of the secondary antibody) and incubated overnight at 4°C with primary antibody diluted at a ratio of 1:1000 in 5% goat or donkey serum in PBS. Secondary antibodies Alexa Fluor 488 or 594 against mouse, rabbit, goat and donkey species were purchased from Life Technologies. Nuclei were visualized by staining with Hoechst 33342 (Sigma-Aldrich). Coverslips were mounted with ProLong® Gold Antifade reagent (Life Technologies) and examined using a Zeiss LSM 710 confocal microscope. Five 25X frames (average of 75 cells each frame) were randomly selected from three independent experiments and regions surrounding

SGs were selected, and corrected total cell fluorescence (CTCF) was calculated using ImageJ. CTCF was used to measure the SG area in arbitrary units also known as brightness values. The number of SGs was manually calculated from five random 63X frames (average of 25 cells each frame) from three independent experiments using Zeiss 2012 software. All statistical calculations were done using GraphPad Prism 6 and one way ANOVA statistical test was performed to calculate the *P*-value. Error bars in the figures represent means \pm SEM.

2.5 **Western blotting and immunoprecipitation assays**

For general immunoblotting experiments, N9, BV-2 or primary microglial cells were lysed in buffer containing 1% NP-40, 50 mM Tris/HCl, pH 8.0, 150 mM NaCl, 5X protease inhibitor cocktail (13911, Sigma-Aldrich), 10 μ M sodium orthovanadate for 15 min on ice. After centrifugation at 14000 X *g* for 10 min, supernatants were collected, separated by SDS-PAGE and analyzed by western blotting. To prepare soluble and insoluble fractions, cells were lysed in buffer A (15 mM Tris-HCl, pH 7.6, 0.3 M NaCl, 15 mM MgCl₂, 1% Triton X-100, 10 mM Ribonucleoside Vanadyl Complex (Cat. No. S1420S, New England Biolabs), 5X protease inhibitor cocktail and 10 μ M sodium orthovanadate) on ice for 10 min. Cells were disrupted by mortar and pestle (20 times moved up and down). The insoluble fraction was isolated by centrifugation at 1500 X *g* for 7 min and the supernatant was collected as the soluble fraction. The insoluble fraction was dissolved in SDS-sample buffer (1% SDS in 1X NP-40 lysis buffer). Further shearing was obtained by passing the cell lysate through a syringe 10 times (28G needle).

Finally western blotting was performed to quantify the protein levels. For phosphotyrosine immunoprecipitation assays, whole cell lysates prepared in buffer A were incubated with anti-phosphotyrosine (4G10, EMD Millipore)-coated protein G magnetic beads (Cat. No. P3296, Sigma-Aldrich) for 2 h at 4°C. Beads were washed thrice with 1X NP-40 lysis buffer, once with 1X high salt solution and once with double-distilled water. Finally, bound proteins were eluted with SDS-sample buffer. Immune complexes were examined by western blotting to identify associated proteins.

GFP-TRAP immunoprecipitation assays were conducted to isolate the GFP-fusion protein from Syk-EGFP-expressing BV-2 cells. Whole cell lysates were prepared in buffer A. A nucleus-enriched portion was separated by centrifuging at 5000 X g for 7 min at 4°C. Supernatant was collected and incubated with GFP-TRAP beads (Cat. No. gta-20, ChromoTek Inc. USA) for 30 min at 4°C. Beads were washed as described above and subjected to western blotting for further analysis.

2.6 Microglial cell functional assays

Phagocytic activity of N9 and BV-2 cells (1×10^4 cells/ plate) was assessed by the uptake of pHrodo™ Red E. coli BioParticles® (Cat. No.P35361, Life Technologies). N9 and BV-2 cells were grown in poly-D-lysine coated glass bottom dishes (MatTek, P35G-1.5-10-C, MatTek Corporation, USA) for 12 h in growth media containing 1% FBS and then treated as indicated for 120 h. Subsequently media was removed. Cells were washed twice with 1X PBS buffer and incubated in 2 ml Live Cell Imaging Solution (Cat. no. A14291DJ, Life Technologies) containing 100 µl pHrodo™ particles (431). Live imaging

was obtained thereafter in 5% CO₂ and 37°C on a confocal microscope (Zeiss LSM 710) stage for another hour. Hoechst dye was added to visualize the nucleus. Live images of cells were taken at 30 sec intervals and compiled into a movie. For fixed cell images, cells were incubated for 1 h in 2 ml Live Cell Imaging Solution containing 100 µl pHrodo™ particles; fixed and examined by confocal microscopy. Phagocytosis of fluorescent red particles was quantified by measuring the mean corrected fluorescence intensity using ImageJ software from five random equal sized frames for each treatment condition.

The phagocytosis of FITC-labeled Aβ fibrils (rPeptide, Cat. No. A-1119-1) was measured using N9 cells in a similar manner except cells were incubated for 1 h with 25 µl FITC-labeled Aβ fibrils (fibrils prepared from 0.25 M solution of soluble Aβ(1-42)). Cells were washed extensively, fixed and imaged by confocal microscopy as mentioned above.

Intracellular ROS production was measured using both the 5(6)-carboxy-2',7'-dichlorodihydrofluorescein diacetate (carboxy-H2DCFDA) (Cat. No. 6982, Setareh Biotech) and dihydrorhodamine 123 (D123) (Cat. No. D-23806, Life Technologies) reagents. N9 cells were grown in 96-well plates in media containing 1% FBS and treated as indicated with SA or Aβ(1-42) in the presence or absence of 500 nM Syk kinase inhibitor (R406 or PRT-060318 (Selleckchem)) for 120 h. Carboxy-H2DCFDA (10 µM) was added and incubated at 37°C for 1 h. Cells were washed 3 times with 1X PBS. Fluorescence was measured using a microplate reader (Biotek Synergy 4). Alternatively, treated and control cells were incubated with D123 (5 µM) for 1 h. Cells were stained with Hoechst dye, fixed and examined by confocal microscopy (Zeiss LSM 710)(91, 432).

Quantitative analyses used ImageJ to measure mean fluorescence intensity. The area enclosing the green fluorescent signal (background signal subtracted) was measured in 5 random frames from three independent experiments.

Extracellular production of H_2O_2 was measured using Acridan Lumigen PS-3 reagents (Amersham ECL kit) as described (433). ALPS-3 substrate was prepared by mixing Reagent A (H_2O_2 , Amersham ECL kit) with Reagent B (Acridan solution, Amersham ECL kit) in a ratio of 10:1. N9 cells grown in media containing 1% FBS were plated on 1% poly-D-lysine coated 24 well plates and treated with SA, $A\beta$ (1-42) and/or Syk inhibitors for 120 h. 50 μ l of media from each well was transferred to another 96 well plate and 50 μ l of fresh ALPS-3 substrate was added. The plate was stored in the dark for 10 min at room temperature. Chemiluminescence was detected using a luminescence plate reader (Glomax, Promega) at 430 nm.

To measure reactive nitrogen species, N9 cells were grown and treated as mentioned above. Media was collected and reactive nitrogen species quantified by measuring nitrite levels using the Measure-iT™ High-Sensitivity Nitrite Assay Kit (Cat. No. M36051, Life technologies). In a black opaque clear bottom 96 well microplate (Corning, USA), 100 μ l of quantitation reagent was incubated with 10 μ l of media from each well containing cells (post-treatment). After incubation for 10 min, 5 μ l of the developing reagent was added and fluorescence was measured using a microplate reader (excitation/emission 365/450 nm, (Biotek Synergy 4)). A standard curve was plotted using the standards provided with the kit.

2.7 Microglia-neuron co-culture and Annexin V assay

N9 microglial cells (5×10^3 cells/well) plated in poly-D-lysine-coated 24-well plates were treated as indicated with SA, A β and/or Syk inhibitor for 120 h. Cells were gently detached and washed with PBS. 5×10^2 cells from each treatment condition were added to an 8 μ m pore size transwell insert (ThincertTM, Greiner bio-one) and placed atop 1.5×10^4 HT22 cells plated in DMEM media without phenol red containing 1% FBS and 100 U/ml penicillin and 100 mg/ml streptomycin. Co-cultured cells were incubated for 48 h. After 48 h, HT22 cells were washed with PBS and incubated with FITC Annexin V (Cat. No. 556420, BD Biosciences) for 30 min at 37°C. Finally cells were fixed, stained with Hoechst dye and examined by confocal microscopy.

2.8 Human AD clinical samples

Paraffin-fixed human AD brain and normal brain cortex samples were obtained from the Brain and Body Donation Program, Banner Sun Health Research Institute, Arizona⁶⁴. Samples were clinically evaluated based on several criteria including mini-mental state examination (MMSE) score, plaque density, lewy body prevalence and dementia. Paraffin fixed cortex brain slices were heated at 65°C for 30 min, fixed with 4% paraformaldehyde, washed sequentially with different concentrations of ethanol (70%, 50%, 30%) and then subjected to antigen retrieval by heating in 10 mM Tris/HCl (pH 10) for 20 min at 85°C. Slides were blocked with 5% donkey serum and 2% BSA overnight at 4°C. Brain slices were stained using primary antibodies against SYK, IBA1, G3BP and phosphotyrosine overnight at 4°C. On the following day, slides were washed 5 times with PBST, stained with Alexa Fluor secondary antibodies, fixed with VectorMount AQ (Cat.

H-5501, Vector Labs) and examined using a Zeiss 710 LSM confocal microscope. The area enclosed by SG in brain slides was measured using ImageJ as mentioned earlier for SGs in cultured cells.

2.9 **Antibodies**

The antibodies and sources used in this study were as follows: Syk (D3Z1E, Cell Signaling Technology or B01P, Abnova), G3BP1 (611126, BD Biosciences or 07-1801, EMD Millipore), pEIF2 α (E90, Abcam), TIA-1(ab2712, Abcam), PABP1 (ab21060, Abcam), GFP (168AT1211, Abgent), GAPDH (6C5, Ambion), pTYR (4G10, EMD Millipore), IBA1 (ab5076, Abcam), GFAP (GF5, Abcam), TDP-43 (10782-2-AP, Proteintech), NOS2 (ab3523, Abcam), Integrin β 1 (4B7R, Santa Cruz Biotech), CD32 (ab197930, Abcam), and Rabbit IgG (Sigma-Aldrich).

CHAPTER THREE: RESULTS

3.1. **SYK is highly expressed in BV-2 and N9 microglia cell**

A few studies have suggested that SYK is present in microglial cells, but none of these compared the expression levels of SYK to those of other immune cells. Endogenous SYK protein level had not been examined previously in N9 cells. Hence, I first wanted to confirm the expression level of SYK in two well-known immortalized microglial cell lines: BV-2 and N9. Cells were grown for 24 h in complete media and lysed in 1X NP-40 lysis buffer. Immunoblotting for endogenous SYK revealed a high level of expression of SYK in both lines as shown in comparison to DG75 B cells (Fig. 3.1). It has been previously observed in our lab that DG175 B cells express substantial levels of SYK .




Figure 3.1 SYK is highly expressed in N9 and BV-2 microglial cell

Figure 3.1 SYK expression was examined in the DG75, B cell-line, and the BV-2 and N9 microglial cell lines. Cell lysates were analyzed by western blotting with SYK and GAPDH antibodies. GAPDH was detected as a loading control.

3.2 **Low dose of sodium arsenite promotes SG formation in microglial cells**

Environmental stress induces SG formation in eukaryotic cells. One of the classical SG inducers is sodium arsenite (SA). To determine if SGs form in these MG, I first treated BV-2 and N9 cells with sodium arsenite for 24 h. Numerous SGs were found in both cell lines when treated with low concentrations of SA (1 μ M) for 24 h as revealed by the appearance of puncta that stained with the SG marker G3BP1 (Fig. 3.2). Next I wanted to examine if SYK localized to these SGs. To determine that, I treated both MG as described above in Fig.3.2, and fixed and stained cells with an antibody against the kinase. A fraction of SYK co-localized with G3BP1 in SGs in both cell types when treated with SA (Fig. 3.2). The concentration of SA that was required to induce SGs was surprisingly low compared to the level commonly (250/500 μ M) used to induce SGs in other cell types (417, 434, 435).




Figure 3.2 SA stimulation manifests SG formation in N9 and BV-2 cells

Figure 3.2 SG formation in BV-2 and N9 cells was examined under SA treatment. BV-2 and N9 cells were treated with SA (1 μ M) for 24 h. On the following day, cells were fixed and stained for endogenous SYK (green) and G3BP1 (red). Confocal images were obtained using a Zeiss LSM 710 microscope. White arrows indicate SYK and G3BP1 co-localized SGs. Scale = 10 μ m.

3.3 **Soluble amyloid beta induces SG formation in MG**

Recent reports have shown that misfolded or aggregated proteins like TAU, TDP-43 and PrP can induce SG formation in neuronal cells (38, 436, 437). Neurodegenerative disease animal models have also confirmed the presence of misfolded proteins associated in SGs in neuronal cells. It is hypothesized that stressors in neurodegeneration can induce such SGs. I then asked if exposure to a stressor implicated in neurodegeneration might also lead to SG formation in MG. Abundant SYK-positive SGs were induced in both BV-2 and N9 cells treated with 100 nM soluble A β (1-42) for 24 h (Fig. 3.3A). Quantitative analyses of SYK and G3BP1 co-localized SG punctas were made for BV- 2 cells using ImageJ. Analyses revealed a five-fold increase in area containing SYK and G3BP1 positive SG in cells treated with SA or A β (1-42) compared to control (Fig. 3.3B). As a control, BV-2 cells also were treated with soluble A β (42-1). A β (42-1) treatment did not form SGs in BV-2 cells (Fig. 3.3C)




Figure 3.3 A β stimulation manifests SG formation in N9 and BV-2 cells

Figure 3.3. (A) BV-2 and N9 cells were treated without (Control) or with A β for 24 h, fixed and stained for SYK (green) and G3BP1 (red). Examples of SGs are indicated by the arrows. (B) The area occupied by SYK and G3BP1 positive SGs in N9 cells was quantified by Image J analysis of 5 random frames from 3 independent experiments. Results represent means \pm SEM. $*P = 0.001$. (C) BV-2 cells were treated with soluble A β (42-1) as a control for 24 h, fixed and stained for SYK (green) and G3BP1 (red). Only merged images are shown. Bar = 10 μ m.

3.4 **SYK and G3BP1 co-localizes with other SG core components in BV-2 and N9 cells**

SGs are extremely dynamic and varied in terms of their components. Various translation initiating factors, mRNA-binding proteins and some other proteins that associate with SG core proteins form the major components of SGs. Depending on the biological system, cell line and stress type, different SG components have been reported. I then asked if other proteins commonly found in SGs were present within these complexes in MG. BV2 cells were treated with SA or A β for 24 h, fixed and stained for SYK and co-stained for phosphorylated EIF2 α , PABP1, TDP-43 or TIA-1. SYK-positive puncta produced by both stimuli contained each of these proteins (Fig. 3.4A). Similar results were observed in N9 cells under similar treatment conditions (Fig. 3.4B). G3BP1 also co-localized in puncta with pEIF2 α , TDP-43 and TIA-1 in N9 cells treated with SA or A β (Fig. 3.5).

Since SGs are insoluble in mild detergents, I fractionated control and stressed MG into detergent-soluble and insoluble fractions and immunoblotted each for SYK, G3BP1, pEIF2 α and TIA-1. A significant increase in the presence of SYK, G3BP1, pEIF2 α and TIA-1 in the detergent-insoluble fractions accompanied treatment of both N9 and BV-2 cells with either SA or A β (Fig. 3.6).




Figure 3.4 SYK co-localizes with pEIF2α, TDP-43, PABP1 and TIA-1 in BV-2 and N9 cells

Figure 3.4. BV-2 (A) and N9 (B) cells were treated without (Control) or with SA or soluble Aβ for 24 h, fixed and stained for SYK, pEIF2α, PABP1, TDP-43, TIA-1 and/or G3BP1 as indicated. Only merged images are displayed. Examples of SGs are indicated by the arrows. Bar = 10 μm.




Figure 3.5. G3BP1 co-localizes with pEIF2α, TDP-43, PABP1 and TIA-1 in BV-2 and N9 cells

Figure 3.5. N9 cells were treated without (Control) or with SA or soluble Aβ for 24 h, fixed and stained for G3BP1, pEIF2α, TDP-43 and TIA-1. Only merged images are displayed. Examples of SGs are indicated by the arrows. Bar = 10 μm.




Figure 3.6 SG markers like SYK, G3BP1, TIA-1 and pEIF2α expression increases in SDS soluble fraction in BV-2 and N9 cells

Figure 3.6 N9 (A) and BV-2 (B) cells were treated without (-) or with (+) SA or soluble Aβ for 24 h and then separated into detergent soluble and insoluble fractions, which were analyzed by western blotting for SYK, G3BP1, TIA-1 and pEIF2α. SA and Aβ showed an increase in protein levels in insoluble fractions under treatment.

3.5 **SYK associates with other SG core components in BV-2 cells**

SG core components associate with other translation initiating factors, mRNA-binding proteins and proteins that interact with core components. To further confirm an interaction or association of SYK with SG markers, I expressed SYK-EGFP in BV-2 cells, which were then treated with SA or A β for 24 hours. Following treatment, cells were lysed and SYK-EGFP was immunoprecipitated using GFP-TRAP beads.

Immunoprecipitated SYK-EGFP was immunoblotted for endogenous SYK, G3BP1 and pEIF2 α . Both stress stimuli resulted in an increased association of SYK-EGFP with endogenous SYK, G3BP1 and pEIF2 α (Fig. 3.7).




Figure 3.7 SG markers G3BP1 and pEIF2 α associate with GFP-tagged SYK in SYK-EGFP-expressing BV-2 cells

Figure 3.7 SYK-EGFP was immunoprecipitated from lentiviral-infected BV-2 cells using GFP-trap beads. Cell lysates (Input) and immune complexes were analyzed by western blotting using antibodies against GFP, SYK, G3BP1 and pEIF2 α . Unmodified beads were used as a control (Mock).

3.6 **SYK is active when present in SGs**

SYK is a protein tyrosine kinase that phosphorylates its substrates to regulate signaling pathways. In immune cells like macrophages and microglia, active SYK mediates downstream signaling that could activate pathways producing reactive oxygen and nitrogen species. To determine if SYK was active when recruited to SGs, I stained control and SA- and A β -treated N9 and BV-2 cells with antibodies against SYK and phosphotyrosine. SYK-positive puncta induced by either stimulus contained abundant phosphotyrosine indicating that SYK was active in SGs (Fig. 3.8A and 3.9).

Quantification of images confirmed a significant increase in area containing co-localized SYK and phosphotyrosine in treated compared to control N9 cells (Fig. 3.8B). Western blotting of lysates of control and SA- or A β -treated N9 cells with antibodies specific for the phosphorylated activation loop of SYK confirmed that SYK was active and phosphorylated in response to both stimuli (Fig. 3.8C). To further confirm the presence of active SYK in SGs biochemically, I immunoprecipitated tyrosine-phosphorylated proteins from control, SA- and A β -treated N9 cells using magnetic beads and immunoblotted the immune complexes for phosphotyrosine and SYK. Induction of SGs with either stimulus resulted in a significant increase in the phosphorylation of multiple proteins including SYK itself (Fig. 3.8D).




Figure 3.8 SYK is active when co-localized in SGs in N9 cells under treatment of SA and Aβ

Figure 3.8. (A) N9 cells were treated without (Control) or with SA or soluble Aβ for 24 h, fixed and stained for SYK (green) and phosphotyrosine (pTYR, red). Examples of SGs are indicated by the arrows. Bar = 10 μm. (B) The area occupied by SYK and phosphotyrosine positive SGs in (A) was quantified by ImageJ analysis of 5 random frames from 3 independent experiments. Results represent means ± SEM. **P* = 0.001. (C) Lysates from N9 cells treated without (Control) or with SA or soluble Aβ for 24 h were analyzed by western blotting using antibodies against SYK phosphorylated on Y519 and Y520 (p-SYK), SYK or GAPDH.

(D) Phosphotyrosine-containing proteins were immunoprecipitated from lysates of N9 cells treated without (Control) or with SA or soluble Aβ for 24 h. Immune complexes were examined by western blotting with antibodies against SYK or phosphotyrosine (pTYR).




Figure 3.9. SYK is active when co-localized in SGs in BV-2 cells under SA and A β treatment

Figure 3.9 BV-2 cells were treated without (Control) or with SA or soluble A β for 24 h, fixed and stained for SYK (green) and phosphotyrosine (pTyr, red). Examples of SGs are indicated by the arrows. Bar = 10 μ m.

3.7 SYK regulates SG formation in MG under SA and A β treatment

SG dynamics are a function of their formation and clearance. To determine if SYK plays an important role in SG formation, I stimulated BV-2 cells with SA or A β for 24 h in the absence or presence of a SYK inhibitor, either R406 or PRT318. In the presence of either inhibitor, neither SA nor A β induced the formation of significant SGs (Fig. 3.10A and 3.10B). The ability of each inhibitor to reduce SYK activity was confirmed by the western blotting with anti-SYK and antiphosphotyrosine antibodies of lysates of BV-2 cells treated with H₂O₂ in the presence or absence of R406 or PRT318. H₂O₂, an inhibitor of tyrosine phosphatases, led to the activation of SYK and its autophosphorylation on Y317 with a resulting shift in its electrophoretic mobility³⁷, which was lost in inhibitor-treated cells (Fig. 3.10C).




Figure 3.10 SG formation dynamics is affected by SYK

Figure 3.10. (A) N9 cells were treated without (Control) or with SA or soluble A β for 24 h in the presence or absence of R406 or PRT318 (PRT), fixed and stained for SYK (green) and G3BP1 (red). Examples of SGs are indicated by the arrows. Bar = 10 μ m. (B) The average number of SGs formed in five random 25X frames were counted for each treatment. Results represent means \pm SEM from 3 independent experiments. * P = 0.0001. (C) N9 cells were treated without (-) or with (+) H₂O₂ (5 μ M) for 4 h in the absence (-) or presence (+) of PRT318 or R406 (500 nM each). Lysates were examined by western blotting for SYK, phosphotyrosine (pTYR) and GAPDH.

3.8 **Knockdown of SYK prevents SG formation in MG under SA and A β treatment**

SYK inhibitors reduced SG formation in N9 cells induced by SA and A β treatment. To confirm a direct role for SYK in SG formation, I infected BV-2 cells with lentiviruses expressing EGFP and either shRNA against SYK or a scrambled shRNA. SYK levels were selectively reduced in cells expressing the SYK shRNA (Fig. 3.11A) as confirmed by western blotting. Infected cells were treated with SA or A β for 24 hours, fixed and stained for G3BP1. While cells infected with scrambled shRNA readily formed SGs, cells with reduced levels of SYK failed to do so (Fig. 3.11B). Correspondingly, the appearance of SYK positive SGs was reduced in the knockdown cells (Fig. 3.12).




Figure 3.11 SYK knockdown prevents G3BP1 associated SG formation in BV-2 cells

Figure 3.11 (A) BV-2 cells expressing EGFP and scrambled shRNA or SYK shRNA were examined by western blotting with antibodies against G3BP1, GFP and GAPDH. (B) BV-2 cells infected with a lentivirus expressing EGFP and either a SYK or scrambled shRNA were treated without (Control) or with SA or soluble A β for 24 h, fixed and stained for G3BP1 (red). Examples of SGs are indicated by the arrows. Bar = 10 μ m.




Figure 3.12 SYK knockdown prevents SYK associated SG formation in BV-2 cells

Figure 3.12 (A) BV-2 cells infected with a lentivirus expressing EGFP and either a SYK or scrambled shRNA were treated without (Control) or with SA or soluble A β for 24 h, fixed and stained for endogenous SYK (red). Examples of SGs are indicated by the arrows. Bar = 10 μ m. (B) Quantification of G3BP1 positive SGs present within 10 random 25X frames for cells expressing SYK shRNA (SYK (-)) or scrambled shRNA (Scram). * P = 0.001.

MG cells lose their efficiency to phagocytose with age. Neurodegeneration is more prevalent in aged individuals. We used a haploinsufficient SYK mouse to conduct these studies. Since SYK knockout is lethal to pups, we had to use haplo-insufficient SYK mice to observe the role of microglial cells in old animal models. In order to first confirm that SYK was indeed knocked down in dead pups or reduced in haploinsufficient mice, we isolated liver cells from the mouse, lysed and immunoblotted for SYK and GAPDH (Fig. 3.13 B). To better understand the role of age on MG, I isolated primary MG from 20 month and 1 month old wild-type mice and haplosufficient mice. Isolated cells were positive for the MG marker IBA-1, but lacked the astrocyte marker GFAP (Fig. 3.13A and C). Cells were treated with SA or A β for 24 h, fixed and stained for IBA-1, SYK and G3BP1. Primary MG formed SYK- and G3BP1-positive puncta in response to both stress stimuli, which were particularly abundant in cells isolated from aged (20 month) animals (Fig. 3.14C). SG formation also was observed, albeit to a lesser extent, in cells isolated from 1 month old animals (Fig. 3.14A).

I also asked if the dependence of SG formation on SYK held true for primary cells. MG isolated from the brains of SYK haploinsufficient mice expressed half as much SYK as cells from wild-type mice (Fig. 3.13A). Since a complete SYK knockout was lethal as seen previously (296), I was unable to test cells from *Syk*^{-/-} animals. SG formation was comparable in cells from one month old *Syk*^{+/+} and *Syk*^{+/-} animals (Fig. 3.14A, 3.15A and 3.16). However SG formation was reduced considerably in cells isolated from the brains of 20 month old *Syk*^{+/-} mice as compared to their wild-type counterparts (Fig. 3.15C and 3.16).




Figure 3.13 Primary microglial culture express IBA-1 and haplosufficient mice express reduced SYK in them

Figure 3.13 (A) Adult primary MG were isolated from 1 month old wild-type (Wt) or *Syk*^{+/-} (Het) mice and examined by western blotting with antibodies against IBA1, SYK, GFAP and GAPDH. (A) Lysates of cells from fetal livers isolated from wild-type (Wt) or *Syk*^{-/-} (KO) mice or from livers of 21 day old wild-type (Wt) or *Syk*^{+/-} mice were analyzed by western blotting for SYK and GAPDH. (B) MG were isolated from 1 month old wild-type (Wt) or *Syk*^{+/-} (Het) mice and examined by immunofluorescence with antibodies against IBA1, Hoechst and GFAP. Bar = 10 μm.




Figure 3.14 Adult primary MG from 20 month old mice generates large SGs

Figure 3.14. (A and B) MG isolated from the brains of 1 month old wild-type mice (A) or 20 month old wild-type mice (B) were treated without (Control) or with SA or soluble Aβ for 24 h, fixed and stained for SYK, IBA1 or G3BP1. Examples of SGs are indicated by the arrows. Bar = 5 μm.




Figure 3.15 Adult primary MG from 20 month old mice *Syk*^{+/-} generate large SGs upon stress induction compared to cells from 1 month old *Syk*^{+/-} mice

Figure 3.15. (A and B) MG isolated from the brains of 1 month old *Syk*^{+/-} mice (A) or 20 month old *Syk*^{+/-} mice (B) were treated without (Control) or with SA or soluble Aβ for 24 h, fixed and stained for SYK, IBA1 or G3BP1. Examples of SGs are indicated by the arrows. Bar = 5 μm.




Figure 3.16 Quantitative comparison of SG formation in MG from young (1m) and old (20m), wild-type (Wt) and Syk haploinsufficient (Het) mice

Figure 3.16 The average number of SGs formed in five random 63X frames for SYK wild-type (1 month and 20 month) and Syk^{+/-} (1 month and 20 month) (Fig. 3.13 & 3.14) for each treatment. Results represent means \pm SEM from 3 independent experiments.

3.10 **Prolonged chronic stress induces large and irreversible SGs in N9 cells**

It has been proposed that exposure to chronic stress leads to the formation of persistent SGs in the brain (29). AD animal model P3106 brains have also shown increased SGs with severity of the disease. To examine this in MG, I exposed N9 cells to low levels of SA or A β for a period of five days to mimic chronic stress. Prolonged stress led to the formation of large SGs that contained both SYK and G3BP1 (Fig. 3.17A). To determine if the formation of these SGs was reversible, I removed the stress and allowed cells to recover for 48 h. While the intensity of staining for G3BP1 declined somewhat, the area within the cell occupied by SGs was largely unchanged (Fig. 3.17A and 3.17C). In contrast, most SGs formed in cells exposed to SA or A β for only 24 h were cleared within 24 h after removal of the stress stimulus (Fig. 3.17B).




Figure 3.17 Prolonged chronic stress stimulation manifests persistent and irreversible SG in N9 cells.

Figure 3.17 (A) N9 cells were treated without (Control) or with SA or soluble Aβ for 5 days. Where indicated, cells were cultured in the absence of SA or Aβ for an additional 48 h. Cells were fixed and stained for SYK (green) and G3BP1 (red). Examples of SGs are indicated by the arrows. Bar = 10 μm. (B) N9 cells were treated for 24 h as indicated and then cultured further in the absence of any treatment for another 24 h. Cells were fixed and stained as in (A). Bar = 10 μm. (C) The area occupied by SYK and G3BP1 positive SGs from (A) was quantified by ImageJ analysis of 5 random frames from 3 independent experiments. Results represent means ± SEM. * $P = 0.0001$.

3.11 **Chronic stress induction leads to SYK-dependent generation of intracellular and extracellular ROS**

Activated MG can generate both reactive oxygen and nitrogen species (ROS and RNS) that can be damaging to neighboring neuronal cells (438). To determine if chronic stress led to constitutive ROS generation, I treated N9 cells with SA or A β for 5 days and then measured ROS production over the course of 1 h by monitoring the oxidation of cell permeant DCFDA. Cells exposed to either SA or A β produced substantially more ROS than control cells (Fig. 3.18A). Similar results were seen when I monitored the conversion of dihydrorhodamine 123 to rhodamine 123 as a measure of ROS production (Fig. 3.18C). Consistent with these results, cells chronically exposed to either SA or A β (1-42) released substantially more ROS into the media than did untreated cells (Fig. 3.18B). ROS production in cells treated with either stress stimulus was largely attenuated in the presence of the SYK inhibitor R406 (500 μ M) or PRT318 (500 μ M) (Fig. 3.18A, B and C).




Figure 3.18 Chronic stress stimulation leads to increased ROS generation in MG cells

Figure 3.18 (A) N9 cells were treated without (Control) or with SA or A β for 5 days in the presence or absence of R406 or PRT318 (500 nM each). Cells were incubated with carboxy-DCFDA (10 μ M) for 1 h and examined using a fluorescence plate reader. Data represent means \pm SEM for 3 independent experiments. ** P = 0.0001. (B) N9 cells were treated as in (A). Media was collected and analyzed for ROS using a chemiluminescence assay. Data represent means \pm SEM for 3 independent experiments. ** P = 0.0001. (C) N9 cells treated as in (A) and were incubated with D123 for 1 h. Green fluorescence was observed using confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Bar = 10 μ m.

3.12 **Chronic stress induction leads to SYK-dependent generation of intracellular and extracellular RNS**

MG cells in AD brain acquire an activated phenotype and secrete both chemokines and NO. The induction of NO in MG is mainly mediated by overexpression of NOS2. I examined changes in the release of RNS in chronically activated MG by measuring nitrite levels in the media. Nitrite release increased three-fold from N9 cells that were chronically stimulated by SA or A β (Fig. 3.19B). Again, levels were reduced significantly if cells were activated in the presence of a SYK inhibitor. Since the formation of RNS in MG is mediated by NOS2 (6), I measured NOS2 levels in control, SA- and A β -treated N9 cells (Fig. 3.19A and B). An increase in the expression of NOS2 was observed by western blotting and immunofluorescence following both stress stimuli. The induction of NOS2 was reduced substantially in cells in which the activity of SYK was inhibited (Fig. 3.19A, B and C).




Figure 3.19. Chronic stress stimulation leads to increased RNS generation by MG cells

Figure 3.19. (A) N9 cells were treated without (Control) or with SA or Aβ for 5 days in the presence or absence of R406 or PRT318 (500 nM each). Following treatment, cells were fixed, immunostained with antibodies against NOS2 and observed using confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Scale = 10 μm. (B) N9 cells were treated as in(A). Media was collected and analyzed for nitrite levels using a fluorescent plate reader. Data represent means ± SEM for 3 independent experiments. **P* = 0.001. (C) N9 cells were treated as in (A) and cells were lysed in NP-40 lysis buffer. Lysate was immunoblotted for NOS2 and GAPDH was blotted as a loading control.

3.13 **Chronic stress induced MG cells mediated neuronal cell death through ROS and RNS secretion**

Since ROS and RNS increased in chronically stressed MG, I asked if this could contribute to neuronal cell death. To test this, I co-cultured HT22 neuronal cells with N9 cells separated by a transwell insert. When chronically SA- and A β -treated N9 cells were cultured with healthy HT22 cells for two days, I observed a significant increase in annexin V binding to HT22 cells (Fig. 3.20A). Negligible annexin V staining was observed when HT22 cells were cultured with N9 cells that were pretreated with SA or A β in the presence of PRT318. To confirm an increase in ROS and RNS in the co-culture media containing activated N9 cells, I measured the levels of each generated over the two day incubation period. A substantial increase in both ROS and RNS was observed when SA- or A β -treated N9 cells were co-cultured with HT22 cells (Fig. 3.20B and C). The inhibition of SYK with either PRT318 or R406 reduced the external ROS and RNS levels significantly.




Figure 3.20. Chronically stressed MG cells induce apoptosis of HT22 neuronal cells in MG-neuron co-cultures.

Figure 3.20. (A) N9 cells were treated without (Control) or with SA or A β for 5 days in the presence or absence of R406 or PRT318 (500 nM each). Following treatment, N9 cells were co-cultured with HT22 for another 48 h separated by a transwell insert. HT22 cells were incubated with FITC Annexin V (green) for 30 min, fixed and imaged using confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Bar = 10 μ m. (B and C) HT22 and N9 cells were co-cultured as in (A). Media was collected from the co-culture experiment and analyzed for levels of ROS (B) or RNS (C) in media using a chemiluminescence assay and fluorescence assay as describe

ed previously (Fig. 3.17B and 3.18B). Data represent means \pm SEM for 3 independent experiments. ** P = 0.0001.

3.14 **Chronic stress induction in BV-2 and N9 cells impairs phagocytosis of pH sensitive *E.coli* Rhodo particles**

An important function of MG that is compromised in aged and AD brains is phagocytosis. To determine if chronic stress negatively affects phagocytotic activity, I treated BV-2 or N9 cells with SA or A β for 5 days and then assayed the cells for the uptake of pH sensitive *E. coli* BioParticles, which fluoresce only following ingestion. Cells were either fixed and stained as shown for BV-2 cells (Fig. 3.21A) or examined by live cell imaging for N9 cells. For both cell types, chronic stress induced by either SA or A β strongly impaired the uptake of the pH Rhodo particles. Phagocytic activity was inhibited as well when cells were treated with either R406 or PRT318 even in the absence of SA or A β supporting an important role for SYK in phagocytosis (Fig. 3.21A and B). Uptake of pH Rhodo particles was blocked by the incubation of cells with cytochalasin D, an inhibitor of phagocytosis (Fig. 3.21C).




Figure 3.21 Chronic SA and A β stimulation and SYK inhibition impairs phagocytosis of E.coli Rhodo particles in BV-2 cells

Figure 3.21. (A) BV-2 cells were treated without (Control) or with SA or soluble A β for 5 days in the presence or absence of R406 or PRT381 (500 nM each) and examined for the uptake of pHRodo particles (red). Cells were fixed and examined by confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Bar = 10 μ m. Similar observations were made using N9 cells examined by time lapse live cell imaging. (B) Quantification of (A) showing change in mean fluorescence measured by Image J from ten random frames in three independent experiments. * P = 0.001, ** P = 0.0001. (C) N9 cells incubated for 4 h without or with cytochalasin D (10 μ M) were examined for the uptake of pHRodo particles (red). Cells were fixed and examined by confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Bar = 10 μ m

3.15 **Chronic stress in SG in MG prevents phagocytosis of β -amyloid fibrils**

An important function of MG is to clear β -amyloid plaques. To examine the effects of stress on this process, I treated N9 cells with SA or A β for 5 days and measured the uptake of fluorescein-labeled A β fibrils. Again, control cells readily phagocytosed A β fibrils whereas chronically stressed cells were unable to do so (Fig. 3.22A and B). Similar to soluble A β , A β fibrils also induced the formation of SGs containing both SYK and G3BP1 when added to either BV-2 or N9 cells (Fig. 3.22C).




Figure 3.22 Chronic SA and A β stimulation impairs phagocytosis of FITC-A β fibrils in N9 cells

Figure 3.22. (A) N9 cells treated without (Control) or with SA or soluble A β for 5 days were incubated with FITC-labeled A β fibrils for 12 h, washed, fixed and examined by confocal microscopy. Hoechst dye (blue) was used to stain the nucleus. Scale= 10 μ m. (B) Quantification of images from (A) by Image J analysis of 5 random frames from 3 independent experiments. Results represent means \pm SEM. $**P = 0.0001$. (C) BV-2 or N9 cells were incubated without (Control) or with A β fibrils for 24 h. Cells were fixed and stained with antibodies against Syk (green) and G3BP1 (red). Only merged images are displayed. Arrows indicate the locations of SGs. Bar = 10 μ m.

3.16 **SYK associated SGs are formed in brains of AD patients**

Our studies on MG indicated a connection between prolonged stress and the development of persistent SGs to which SYK was recruited. To determine if a similar mechanism was operative in AD, I examined brain samples from patients with no, mild, moderate or severe AD (Table 1). Paraffin-fixed sections of brain cortex were stained with antibodies against IBA1, SYK, G3BP1 and/or phosphotyrosine. SYK was diffusely distributed in MG in patients with no AD and in patients with mild disease, but was increasingly present in large puncta with escalating disease severity (Fig. 3.23). SYK co-localized within these puncta in diseased brain with both G3BP1 and phosphotyrosine indicating that the kinase was localized to SGs and was active at these sites (Fig. 3.24 and 3.25).

Table 1. Characteristics of Alzheimer's disease patients used to measure SYK, G3BP1 and phosphotyrosine (pTYR) co-localized in SGs.

Disease State	No. of patients	Avg . Age	Avg. MMS E score	Total plaques	Total tangles	Area covered by SG Puncta (SYK & G3BP1) Fluorescence	Area covered by SG Puncta (SYK & pTYR) Fluorescence
Normal	6	77	30	3	2	148	234
Mild	6	91	26	1.5	4	268	435
Moderate	6	91	22	7	5	675	746
Severe	6	82	10	14	14	1357*	1467*

The mean area occupied by SGs positive for SYK and G3BP1 or SYK and pTYR was quantified by Image J analysis of 5 random frames from 6 different patient slides of each disease state. * $P = 0.001$.




Figure 3.23. SYK and G3BP1 positive granules were observed in human AD brain samples

Figure 3.23 (A and B) Paraffin embedded slides of human brain cortex from patients with no (Normal), mild, moderate (Mod) and severe AD were immunostained for IBA1 (microglia marker), SYK and G3BP1 independently as indicated. Bar = 5 μ m.




Figure 3.24 SYK and G3BP1 positive granules co-localized in human AD brain samples

Figure 3.24 Paraffin embedded slides of human brain cortex from patients with no (Normal), mild, moderate (Mod) and severe AD were immunostained for SYK and G3BP1 as indicated. Bar = 5 μ m.




Figure 3.25 SYK positive granules co-localized with phosphotyrosine in human AD brain samples

Figure 3.25 Paraffin embedded slides of human brain cortex from patients with no (Normal), mild, moderate (Mod) and severe AD were immunostained for SYK and phosphotyrosine as indicated. Bar = 5 μ m.

3.17 **IgG treatment can repair impaired phagocytosis of MG after chronic stress induction.**

Since phagocytosis of MG was impaired under SA or A β stimulation, I wanted to devise a way for cells to recover from phagocytosis impairment. I asked if manipulations of SYK activity or localization might restore the ability of chronically stressed MG to phagocytose foreign particles. Since integrin ligation leads to the activation of SYK in macrophages (439), I first treated chronically stressed N9 cells with rabbit IgG directed against β 1 integrin. Treatment of stressed N9 cells for 4 h led to a marked recovery of phagocytic activity (Fig. 3.25). As a control, I treated stressed N9 cells with nonspecific rabbit IgG isotype control antibodies. Interestingly, phagocytic activity again was restored. To explore this further, I treated stressed cells with an affinity purified rabbit IgG directed against a peptide from the cytoplasmic C-terminus of the inhibitory receptor CD32, an antibody that should neither react with intact MG nor activate SYK. Again, incubation with this antibody led to a recovery of phagocytic activity in chronically stressed cells (Fig. 3.25). Thus, phagocytic activity in stressed MG could be restored by treatment with IgG independent of the specificity of the antibody.

To explore the mechanism further, I examined the effects of IgG on the localization of SYK. Treatment of N9 cells with either non-specific rabbit IgG or the anti-CD32 antibody led to a relocalization of SYK from the cytoplasm to the plasma membrane, the site where phagocytosis takes place. Even in chronically stressed cells, a substantial amount of SYK was translocated from SGs to the membrane (Fig. 3.26). Western blotting

of detergent-soluble and -insoluble fractions of N9 cells confirmed the partial re-location of SYK from the insoluble fraction in which SGs are located to the detergent-soluble fraction following incubation with rabbit IgG or anti-CD32 (Fig. 3.27A and B).




Figure 3.26 IgG treatment of chronically stressed N9 cells restores phagocytic activity.

Figure 3.26. N9 cells were treated without (Control) or with SA or soluble A β for 120 h. At 116 h, to one set of control, SA- and A β -treated cells were treated with anti- β 1-integrin (β -Int), mixed rabbit IgG (IgG) or anti-CD32 for the final 4 h. Cells were then incubated with pHRhodo particles (red) to measure phagocytosis. Cells were fixed and stained with Hoechst dye (blue) to visualize the nucleus. Bar = 10 μ m.




Figure 3.27. IgG treatment of chronically stressed N9 cells relocates SYK to the plasma membrane

Figure 3.27. N9 cells were treated without (Control) or with SA or soluble A β for 120 h in the presence or absence of R406 or PRT318 (500 nM each). At 116 h, to one set of control, SA- and A β -treated cells were treated with mixed rabbit IgG (IgG) or anti-CD32 for the final 4 h. Cells were then fixed and stained with antibodies against endogenous SYK. Bar = 10 μ m.




Figure 3.28 IgG treatment of chronically stressed N9 cells partially relocalizes SYK from the SG insoluble fraction to the soluble fraction.

Figure 3.27 (A and B) N9 cells were treated without (Control) or with SA or soluble A β for 120 h. At 116 h, one set of control, SA- and A β -treated cells were treated with anti-CD32 (A) or rabbit IgG (Rab IgG) (B) for the final 4 h. Cells were separated into detergent soluble and insoluble fractions, which were analyzed by western blotting for SYK and GAPDH.

CHAPTER FOUR: DISCUSSION

Paradoxically, MG from aged and AD brain exhibit a phenotype that combines a perpetually activated state with functionally impaired phagocytic activity (3, 6). To explain this dichotomy, we propose a model in which the altered activity and localization of SYK contributes to both phenotypes. In this model, chronic stress leads to the formation of SGs to which SYK is recruited resulting both in a persistent pro-inflammatory phenotype and impaired phagocytosis.

SG formation is thought to be, at least initially, a defense against stress, providing a mechanism for the selective sequestration and protection from degradation of a subset of mRNAs (27). However, it is clear that persistent formation of RNP particles is deleterious to cells as noted for several neurodegenerative disorders where aggregates appear in neuronal cells (31, 40). Our data indicate that SGs form, as well, in MG. In fact, MG appears particularly susceptible to the stress-induced formation of SGs as their induction requires exposure only to low levels of SA. The recruitment of SYK to SGs in non-inflammatory cells protects them from stress-induced damage by stimulating SG clearance through autophagy, an event that requires both SYK activity and its association with SGs (Krisenko et al., manuscript submitted for publication). This process may be operative as well in MG as SGs can be cleared effectively from MG exposed to stress for short periods of time. However, prolonged stress leads to the formation of large SGs that

associate with SYK and are resistant to clearance as they persist following removal of the stimulus. SYK in SGs is active as reflected by the presence of phosphotyrosine-containing proteins, which is a function of SYK's catalytic activity (Krisenko et al., manuscript submitted for publication). Large SGs also are characteristic of MG in the brains of patients with advanced AD, and these also contain abundant SYK and phosphotyrosine. Thus, the appearance of SYK- and phosphotyrosine-containing puncta in brain samples may serve as a useful marker of dysfunctional MG in neurodegenerative disease.




Figure 4.1 Model of SYK associated stress granules formation in microglial cell

SA induces SG formation in part through the generation of ROS⁴¹. Oxidative stress also is thought to precede the pathogenesis of AD since enhanced neuroinflammation resulting from traumatic brain injury, infection, or aging predispose individuals to AD while the long-term use of NSAIDS can be protective (440). We find that when stress stimulates the formation of SGs in MG this hampers their ability to phagocytose A β (1-42) fibrils. Thus, we predict that enhanced accumulation of A β plaques in the brain would be an expected result of a prolonged exposure to oxidative stress. We find further that the exposure of MG to A β or A β fibrils alone also induces the appearance of abundant SGs. This is consistent with the known ability of A β to trigger MG cell activation and promote the generation of ROS and RNS (9, 376). Thus, a defect in the removal of A β fibrils by MG compromised by oxidative stress could cause a build-up of A β plaques leading to additional A β -induced stress and the enhanced formation of persistent SGs. MG from aged mice are more sensitive to the stress-induced formation of SGs suggesting one mechanism by which age might lead to a greater susceptibility to AD.




Figure 4.2 Graph art depicting stress granule formation in microglial cell upon Aβ stimulation

The formation of SGs in MG in response to low levels of either SA or A β requires the expression of SYK as it is blocked by SYK inhibitors or by a knockdown or reduction of SYK expression. In inflammatory cells like MG, SYK plays a well known role in the receptor-mediated production of ROS and RNS (271). A β binds to MG via an assortment of cell surface receptors (6) many of which, including CD14, TLR4, CD36, and SIRP β 1, are coupled to SYK either directly or through receptor-associated proteins that contain ITAMs (254, 367, 383, 441, 442). SYK also is activated in cells under conditions of increased oxidative stress (443). The resulting SYK-dependent pro-inflammatory response may be further exacerbated by the retention within SGs of active SYK, which may continue to stimulate down-stream signaling pathways that generate ROS or RNS. The production of ROS and RNS can be damaging to neighboring cells (6) and we find, in fact, that chronically activated MG produce mediators that are toxic to HT22 neuronal cells. Consistent with a role for SYK in these processes, the generation of chronically activated MG that produce ROS and RNS and the ability of these cells to kill neuronal cells are all blocked by SYK inhibitors.

The ability of MG to clear A β plaques through phagocytosis is important for the control of AD (5, 439). We find that chronic stress induced by prolonged exposure of MG to either SA or A β inhibits the ability of these cells to phagocytose either bacteria or A β fibrils. Interestingly, in a mouse model of AD, the phagocytic activity of MG also is selectively compromised in brain regions containing A β plaques (4). We find that chronic stimulation of MG with either SA or A β results in the formation of very large SGs to which SYK is recruited and propose that it is the sequestration of the kinase away from

phagocytic receptors that underlies the impaired phagocytic activity of MG encountering A β plaques.

A role for SYK in both phagocytosis and the activation of MG complicates the development of strategies for the use of specific SYK inhibitors for the treatment or prevention of AD. However, the finding that active SYK is sequestered in SGs suggests that strategies to relocate the enzyme might be effective in restoring some function to damaged MG. We find that the treatment of chronically stressed MG with rabbit IgG leads to a dramatic restoration of phagocytic activity. The recovery of activity is not a function of the reactivity of the IgG to any specific antigen as even an affinity-purified antibody prepared against a peptide found on the cytoplasmic domain of an ITIM-containing receptor is able to induce this effect. Treatment of MG with IgG leads to a relocalization of SYK from the cytoplasm to the plasma membrane in unstressed cells and from SGs to the membrane in chronically stressed cells. This effect may help explain the therapeutic benefits of intravenous immunoglobulins (IVIg), which have shown efficacy in slowing the progression of AD in some human clinical trials (444-446). The benefits of IVIg are generally attributed to the presence of anti-A β antibodies in the pooled IgG preparations (444). However, it is interesting to note that the administration of pooled mouse IgG having no detectable anti-A β activity to the brains of APP/PS1 mice leads to a similar reduction in A β deposits as does IVIg (447). Thus, even immunoglobulins that fail to recognize A β have the potential to reverse defects in the function of stressed MG. This observation suggests that the development of IgG-related therapeutics optimized for the recovery of phagocytic activity of MG would be an attractive strategy for augmenting current strategies for the treatment of AD.

CHAPTER FIVE: FUTURE DIRECTIONS

Stress granule formation in neurodegenerative brain has been observed in several diseases including ALS, AD, SMA and FXMR(29). Autopsy studies on the brains of these patients have indicated an increased aggregation of SG-associated proteins with increasing severity of the disease. Our studies showed for the first time the presence of SGs in MG. MG over-activation is the leading cause of neuroinflammation in neurodegenerative diseases. Our data show that SYK promotes SG formation in MG under SA or A β mediated stress. It would be very interesting to further dissect the other stressors that influence the formation of SGs in MG. We know, for example, that several genetic mutations in genes like *tau*, *PINK1*, *PARKIN*, *APP*, *SNCA*, *CDK5*, *MST1* and *prion* directly manifests disease symptoms (448, 449). To better understand the role of SGs in these diseases, it would be interesting to examine these mutant proteins to determine if they directly or indirectly drive SG formation or inhibit SG clearance in the brain during disease progression.

In this study, we discovered that chronic stress at a very low dose can induce SGs that are persistent and cannot be removed upon removal of stress. We also observed that SYK localizes in SGs and influences their formation in these cells. Additionally, SYK regulates ROS and RNS production of MG cells. Many pathways dissected in AD have

indicated that excessive production of ROS and RNS hastens neuronal cell death through chronic neuroinflammation. Our study also showed that SYK associates with other misfolded proteins like TDP-43 upon stress stimulation. Syk was also observed to be active in these SGs. SYK is a tyrosine kinase that phosphorylates several downstream proteins to trigger activation of other pathways such as Ras/ERK and activate transcription factors such as NF- κ B (138, 224, 450). These downstream proteins directly or indirectly modulate release of pro-inflammatory factors, cytokines or reactive species in various cells including immune cells. A proteomic screen of MG under acute or chronic stress will help to identify the proteins that associate with SYK in SGs under chronic stress. Furthermore, we can also identify the major pathways that are activated that result in SG formation. It is well known that SYK associates with ITAM bearing receptor proteins in MG. A detailed tyrosine phosphoproteomic screen could identify the receptors that activate SYK under SG formation conditions. Several MG receptors like Fc γ R, integrins, CD33 and CD-3 are involved in engulfing A β plaques and tau tangles in AD (341, 451). The screening could also reveal the receptors that lose their ability to engulf foreign particles like misfolded proteins under chronic stress.

Autophagy is a mechanism by which cells get rid of intracellular protein aggregates and damaged organelles. The intracellular proteins are engulfed by double walled vacuoles and delivered to lysosomes for degradation. In neurodegenerative diseases, studies have shown that autophagy mediates removal of impaired organelles and toxic protein aggregates like HTT, tau and beta amyloid. Impairment of autophagy is also observed in neurodegenerative diseases like AD, PD and HD (32, 410). Additionally, several studies suggest that promoting autophagy alleviates disease symptoms in AD and

HD. On the other hand, it has been reported in a few studies that autophagy could be one of the mechanisms by which SGs are cleared from the cell. Based on such evidence, it could be proposed that impairment of autophagy leads to an increase in the number of pathological SGs in neurodegenerative brain. Due to reduced autophagy, a greater number of toxic protein aggregates are sequestered to the SGs and a chain of downstream unfavorable reactions occur. Understanding the connection between autophagy and SGs in neurodegenerative brain, specifically MG, could provide us with better therapeutic approaches to retarding disease progression. It would be also interesting to examine if promoting autophagy would increase clearance of pathological SGs resulting in alleviated disease symptoms.

One of the vital functions of MG is phagocytosis of protein aggregates, debris and dead cells. MG efficiently perform phagocytosis of amyloid plaques and tau fibrils in early AD brain, but fail to do so with disease progression. Additionally MG adopt an activated state, releasing cytokines, chemokines and reactive species contributing to inflammation (124, 430). In our study, we have shown that inhibition of SYK in A β stressed MG can prevent the generation of such reactive species. Sadly, SYK inhibition also impairs phagocytosis. But another finding in our study also showed that IgG treatment can re-localize SYK partially to the plasma membrane from SGs. This re-localization reconstituted the phagocytosis machinery in MG and the number of SGs also was reduced. Further studies need to be done to dissect which receptors IgG binds to on the membrane to provoke such a relocalization of SYK. It would be interesting to determine if the binding of IgG Fc regions to the receptors is enough to relocalize SYK and if specific Fc γ receptors are required for this process. A knowledge of IgG binding

and SYK relocalization could prove to be a promising therapeutic approach for repairing MG in AD.

Since this study led to some interesting findings about SYK's role in MG activation in AD; it would be logical to expand this study to neurodegenerative animal models. APP/PS1 is a very suitable mouse transgenic model to study A β mediated neurodegeneration in AD (131, 222). Our study suggests that SYK inhibition and IVIg can be a promising therapeutic approach to repair impaired MG in AD. An animal model study involving use of SYK inhibitors and IVIg independently or in combination in APP/PS1 transgenic mice would be interesting. Examination of neuroinflammation and MG activation in those animal models will tell us more about microglial SYK's role in AD. Additionally, behavioral studies would also help us determine the importance of this study in the development of therapeutics.

Primary MG when stressed with SA or A β developed large SYK and G3BP1 positive SGs. MG from 20 month old mice showed large SGs compared to 1 month old mice. Further examination in SYK haplosufficient mice showed reduced SGs in stressed MG confirming SYK's role in SG formation. Since the SYK knockout mouse is biologically lethal, we are trying to develop a transgenic mouse that expresses an analog sensitive form of SYK (296). Treatment with an orthogonal inhibitor like NM-PP1 can inhibit SYK's activity selectively in these mice at any point in their life cycle (452). It would be interesting to examine the SG dynamics in MG of these mice upon A β stress induction. Subsequent investigation of MG in this genetically modified SYK mouse could give us insights into receptors that are impaired upon SYK knockdown in the long run. We could also detect various behavioral or pathological implications of SYK

knockdown in an adult mouse. The next step would be to develop a transgenic mouse by genetically crossing analog sensitive form of SYK knockout mouse and APP/PS1 mice. Such transgenic mice can further help us to dissect the exact role of SYK in overall progression of AD. By turning off SYK at any specific time point in the life cycle of such transgenic mice, we can identify the role of microglial SYK in early and late AD. We can also determine the various SYK associated ITAM receptors that are activated and impaired at different stages of AD.

LIST OF REFERENCES

LIST OF REFERENCES

1. Huang Y, Mucke L. Alzheimer Mechanisms and Therapeutic Strategies. *Cell* 2012 2012/03;148(6):1204-22.
2. Solito E, Sastre M. Microglia Function in Alzheimer's Disease. *Front Pharmacol* 2012;3.
3. Gandy S, Heppner Frank L. Microglia as Dynamic and Essential Components of the Amyloid Hypothesis. *Neuron* 2013 2013/05;78(4):575-7.
4. Krabbe G, Halle A, Matyash V, Rinnenthal JL, Eom GD, Bernhardt U, Miller KR, Prokop S, Kettenmann H, Heppner FL. Functional Impairment of Microglia Coincides with Beta-Amyloid Deposition in Mice with Alzheimer-Like Pathology. *PLoS One* 2013 2013/04/08;8(4):e60921.
5. Hickman SE, Allison EK, El Khoury J. Microglial dysfunction and defective beta-amyloid clearance pathways in aging Alzheimer's disease mice. *J Neurosci* 2008 Aug 13;28(33):8354-60.
6. Mosher KI, Wyss-Coray T. Microglial dysfunction in brain aging and Alzheimer's disease. *Biochem Pharmacol* 2014 Apr 15;88(4):594-604.
7. Meyer-Luehmann M, Spires-Jones TL, Prada C, Garcia-Alloza M, de Calignon A, Rozkalne A, Koenigsknecht-Talboo J, Holtzman DM, Bacskai BJ, Hyman BT. Rapid appearance and local toxicity of amyloid-beta plaques in a mouse model of Alzheimer's disease. *Nature* 2008 Feb 7;451(7179):720-4.
8. Jekabsone A, Mander PK, Tickler A, Sharpe M, Brown GC. Fibrillar beta-amyloid peptide Abeta1-40 activates microglial proliferation via stimulating TNF-alpha release and H2O2 derived from NADPH oxidase: a cell culture study. *J Neuroinflammation* 2006;3:24.
9. Bianca VD, Dusi S, Bianchini E, Dal Pra I, Rossi F. beta-amyloid activates the O-2 forming NADPH oxidase in microglia, monocytes, and neutrophils. A possible inflammatory mechanism of neuronal damage in Alzheimer's disease. *J Biol Chem* 1999 May 28;274(22):15493-9.
10. Guerreiro R, Wojtas A, Bras J, Carrasquillo M, Rogaeva E, Majounie E, Cruchaga C, Sassi C, Kauwe JSK, Younkin S, Hazrati L, Collinge J, Pocock J, Lashley T, Williams J, Lambert J-C, Amouyel P, Goate A, Rademakers R, Morgan K, Powell J, St. George-Hyslop P, Singleton A, Hardy J. TREM2 Variants in Alzheimer's Disease. *New England Journal of Medicine* 2013 2013/01/10;368(2):117-27.

11. Jonsson T, Stefansson H, Steinberg S, Jonsdottir I, Jonsson PV, Snaedal J, Bjornsson S, Huttenlocher J, Levey AI, Lah JJ, Rujescu D, Hampel H, Giegling I, Andreassen OA, Engedal K, Ulstein I, Djurovic S, Ibrahim-Verbaas C, Hofman A, Ikram MA, van Duijn CM, Thorsteinsdottir U, Kong A, Stefansson K. Variant of TREM2 Associated with the Risk of Alzheimer's Disease. *New England Journal of Medicine* 2013 2013/01/10;368(2):107-16.
12. Bertram L, Lange C, Mullin K, Parkinson M, Hsiao M, Hogan MF, Schjeide BMM, Hooli B, DiVito J, Ionita I, Jiang H, Laird N, Moscarillo T, Ohlsen KL, Elliott K, Wang X, Hu-Lince D, Ryder M, Murphy A, Wagner SL, Blacker D, Becker KD, Tanzi RE. Genome-wide Association Analysis Reveals Putative Alzheimer's Disease Susceptibility Loci in Addition to APOE. *The American Journal of Human Genetics* 2008 2008/11;83(5):623-32.
13. Hollingworth P, Harold D, Sims R, Gerrish A, Lambert J-C, Carrasquillo MM, Abraham R, Hamshere ML, Pahwa JS, Moskvina V, Dowzell K, Jones N, Stretton A, Thomas C, Richards A, Ivanov D, Widdowson C, Chapman J, Lovestone S, Powell J, Proitsi P, Lupton MK, Brayne C, Rubinsztein DC, Gill M, Lawlor B, Lynch A, Brown KS, Passmore PA, Craig D, McGuinness B, Todd S, Holmes C, Mann D, Smith AD, Beaumont H, Warden D, Wilcock G, Love S, Kehoe PG, Hooper NM, Vardy ERLC, Hardy J, Mead S, Fox NC, Rossor M, Collinge J, Maier W, Jessen F, R  ther E, Sch  rmann B, Heun R, K  lsch H, van den Bussche H, Heuser I, Kornhuber J, Wiltfang J, Dichgans M, Fr  lich L, Hampel H, Gallacher J, H  ll M, Rujescu D, Giegling I, Goate AM, Kauwe JSK, Cruchaga C, Nowotny P, Morris JC, Mayo K, Sleegers K, Bettens K, Engelborghs S, De Deyn PP, Van Broeckhoven C, Livingston G, Bass NJ, Gurling H, McQuillin A, Gwilliam R, Deloukas P, Al-Chalabi A, Shaw CE, Tsolaki M, Singleton AB, Guerreiro R, M  hleisen TW, N  then MM, Moebus S, J  ckel K-H, Klopp N, Wichmann HE, Pankratz VS, Sando SB, Aasly JO, Barcikowska M, Wszolek ZK, Dickson DW, Graff-Radford NR, Petersen RC, van Duijn CM, Breteler MMB, Ikram MA, DeStefano AL, Fitzpatrick AL, Lopez O, Launer LJ, Seshadri S, Berr C, Campion D, Epelbaum J, Dartigues J-F, Tzourio C, Alperovitch A, Lathrop M, Feulner TM, Friedrich P, Riehle C, Krawczak M, Schreiber S, Mayhaus M, Nicolhaus S, Wagenpfeil S, Steinberg S, Stefansson H, Stefansson K, Snaedal J, Bj  rnsson S, Jonsson PV, Chouraki V, Genier-Boley B, Hiltunen M, Soininen H, Combarros O, Zelenika D, Delepine M, Bullido MJ, Pasquier F, Mateo I, Frank-Garcia A, Porcellini E, Hanon O, Coto E, Alvarez V, Bosco P, Siciliano G, Mancuso M, Panza F, Solfrizzi V, Nacmias B, Sorbi S, Boss   P, Piccardi P, Arosio B, Annoni G, Seripa D, Pilotto A, Scarpini E, Galimberti D, Brice A, Hannequin D, Licastro F, Jones L, Holmans PA, Jonsson T, Riemenschneider M, Morgan K, Younkin SG, Owen MJ, O'Donovan M, Amouyel P, Williams J. Common variants at ABCA7, MS4A6A/MS4A4E, EPHA1, CD33 and CD2AP are associated with Alzheimer's disease. *Nat Genet* 2011 2011/04/03;43(5):429-35.
14. Naj A, Jun G, Buross J, Gallins P, Farrer L, Haines J, Pericak-Vance M, Schellenberg G. Genome-Wide Association Study of Late-Onset Alzheimer Disease Identifies Disease-Associated Variants in MS4A4/MS4A6E, CD2AP, CD33, and EPHA1. *Alzheimer's & Dementia* 2011 2011/07;7(4):S191.

15. Zhang J, Okabe K, Tani T, Funatsu T. Dynamic association-dissociation and harboring of endogenous mRNAs in stress granules. *J Cell Sci*2011 Dec 1;124(Pt 23):4087-95.
16. N'Diaye EN, Branda CS, Branda SS, Nevarez L, Colonna M, Lowell C, Hamerman JA, Seaman WE. TREM-2 (triggering receptor expressed on myeloid cells 2) is a phagocytic receptor for bacteria. *J Cell Biol*2009 Jan 26;184(2):215-23.
17. Takahashi K. Clearance of apoptotic neurons without inflammation by microglial triggering receptor expressed on myeloid cells-2. *Journal of Experimental Medicine*2005 2005/02/14;201(4):647-57.
18. Wang Y, Cella M, Mallinson K, Ulrich JD, Young KL, Robinette ML, Gilfillan S, Krishnan GM, Sudhakar S, Zinselmeyer BH, Holtzman DM, Cirrito JR, Colonna M. TREM2 Lipid Sensing Sustains the Microglial Response in an Alzheimer's Disease Model. *Cell*2015 Mar 12;160(6):1061-71.
19. Jay TR, Miller CM, Cheng PJ, Graham LC, Bemiller S, Broihier ML, Xu G, Margevicius D, Karlo JC, Sousa GL, Cotleur AC, Butovsky O, Bekris L, Staugaitis SM, Leverenz JB, Pimplikar SW, Landreth GE, Howell GR, Ransohoff RM, Lamb BT. TREM2 deficiency eliminates TREM2+ inflammatory macrophages and ameliorates pathology in Alzheimer's disease mouse models. *J Exp Med*2015 Mar 9;212(3):287-95.
20. Griciuc A, Serrano-Pozo A, Parrado Antonio R, Lesinski Andrea N, Asselin Caroline N, Mullin K, Hooli B, Choi Se H, Hyman Bradley T, Tanzi Rudolph E. Alzheimer's Disease Risk Gene CD33 Inhibits Microglial Uptake of Amyloid Beta. *Neuron*2013 2013/05;78(4):631-43.
21. Linnartz B, Wang Y, Neumann H. Microglial immunoreceptor tyrosine-based activation and inhibition motif signaling in neuroinflammation. *Int J Alzheimers Dis*2010;2010.
22. Cambier JC. New nomenclature for the Reth motif (or ARH1/TAM/ARAM/YXXL). *Immunology Today*1995 1995/01;16(2):110.
23. Geahlen RL. Syk and pTyr'd: Signaling through the B cell antigen receptor. *Biochim Biophys Acta*2009 Jul;1793(7):1115-27.
24. Angata T, Kerr SC, Greaves DR, Varki NM, Crocker PR, Varki A. Cloning and Characterization of Human Siglec-11: A RECENTLY EVOLVED SIGNALING MOLECULE THAT CAN INTERACT WITH SHP-1 AND SHP-2 AND IS EXPRESSED BY TISSUE MACROPHAGES, INCLUDING BRAIN MICROGLIA. *Journal of Biological Chemistry*2002 2002/05/01;277(27):24466-74.
25. Kiefer F, Brumell J, Al-Alawi N, Latour S, Cheng A, Veillette A, Grinstein S, Pawson T. The Syk protein tyrosine kinase is essential for Fcgamma receptor signaling in macrophages and neutrophils. *Mol Cell Biol*1998 Jul;18(7):4209-20.
26. Crowley MT, Costello PS, Fitzer-Attas CJ, Turner M, Meng F, Lowell C, Tybulewicz VL, DeFranco AL. A critical role for Syk in signal transduction and phagocytosis mediated by Fcgamma receptors on macrophages. *J Exp Med*1997 Oct 6;186(7):1027-39.
27. Anderson P, Kedersha N. Stress granules: the Tao of RNA triage. *Trends Biochem Sci*2008 Mar;33(3):141-50.
28. Buchan JR, Parker R. Eukaryotic stress granules: the ins and outs of translation. *Mol Cell*2009 Dec 25;36(6):932-41.

29. Ash PE, Vanderweyde TE, Youmans KL, Apicco DJ, Wolozin B. Pathological stress granules in Alzheimer's disease. *Brain Res*2014 Oct 10;1584:52-8.
30. King OD, Gitler AD, Shorter J. The tip of the iceberg: RNA-binding proteins with prion-like domains in neurodegenerative disease. *Brain Res*2012 Jun 26;1462:61-80.
31. Harris H, Rubinsztein DC. Control of autophagy as a therapy for neurodegenerative disease. *Nat Rev Neurol*2012 Feb;8(2):108-17.
32. Buchan JR, Kolaitis RM, Taylor JP, Parker R. Eukaryotic stress granules are cleared by autophagy and Cdc48/VCP function. *Cell*2013 Jun 20;153(7):1461-74.
33. Koppers M, van Blitterswijk MM, Vlam L, Rowicka PA, van Vught PWJ, Groen EIJ, Spliet WGM, Engelen-Lee J, Schelhaas HJ, de Visser M, van der Kooij AJ, van der Pol WL, Pasterkamp RJ, Veldink JH, van den Berg LH. VCP mutations in familial and sporadic amyotrophic lateral sclerosis. *Neurobiology of Aging*2012 2012/04;33(4):837.e7-.e13.
34. King OD, Gitler AD, Shorter J. The tip of the iceberg: RNA-binding proteins with prion-like domains in neurodegenerative disease. *Brain Research*2012 2012/06;1462:61-80.
35. Vanderweyde T, Yu H, Varnum M, Liu-Yesucevitz L, Citro A, Ikezu T, Duff K, Wolozin B. Contrasting Pathology of the Stress Granule Proteins TIA-1 and G3BP in Tauopathies. *Journal of Neuroscience*2012 2012/06/13;32(24):8270-83.
36. Waelter S, Boeddrich A, Lurz R, Scherzinger E, Lueder G, Lehrach H, Wanker EE. Accumulation of Mutant Huntingtin Fragments in Aggresome-like Inclusion Bodies as a Result of Insufficient Protein Degradation. *Molecular Biology of the Cell*2001 2001/05/01;12(5):1393-407.
37. Goggin K, Beaudoin S, Grenier C, Brown A-A, Roucou X. Prion protein aggresomes are poly(A)+ ribonucleoprotein complexes that induce a PKR-mediated deficient cell stress response. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research*2008 2008/03;1783(3):479-91.
38. Liu-Yesucevitz L, Bilgutay A, Zhang Y-J, Vanderwyde T, Citro A, Mehta T, Zaarur N, McKee A, Bowser R, Sherman M, Petrucelli L, Wolozin B. Tar DNA Binding Protein-43 (TDP-43) Associates with Stress Granules: Analysis of Cultured Cells and Pathological Brain Tissue. *PLoS One*2010 2010/10/11;5(10):e13250.
39. Neumann M, Sampathu DM, Kwong LK, Truax AC, Micsenyi MC, Chou TT, Bruce J, Schuck T, Grossman M, Clark CM, McCluskey LF, Miller BL, Masliah E, Mackenzie IR, Feldman H, Feiden W, Kretschmar HA, Trojanowski JQ, Lee VMY. Ubiquitinated TDP-43 in Frontotemporal Lobar Degeneration and Amyotrophic Lateral Sclerosis. *Science*2006 2006/10/06;314(5796):130-3.
40. Menzies FM, Fleming A, Rubinsztein DC. Compromised autophagy and neurodegenerative diseases. *Nat Rev Neurosci.* [Review]. 2015;16(6):345-57.
41. Iliuk AB, Martin VA, Alicie BM, Geahlen RL, Tao WA. In-depth Analyses of Kinase-dependent Tyrosine Phosphoproteomes Based on Metal Ion-functionalized Soluble Nanopolymers. *Molecular & Cellular Proteomics*2010 2010/06/17;9(10):2162-72.
42. Xue L, Wang WH, Iliuk A, Hu L, Galan JA, Yu S, Hans M, Geahlen RL, Tao WA. Sensitive kinase assay linked with phosphoproteomics for identifying direct kinase substrates. *Proceedings of the National Academy of Sciences*2012 2012/03/26;109(15):5615-20.

43. Galan JA, Paris LL, Zhang H-j, Adler J, Geahlen RL, Tao WA. Proteomic Studies of Syk-Interacting Proteins Using a Novel Amine-Specific Isotope Tag and GFP Nanotrap. *J Am Soc Mass Spectrom* 2011 2011/01/28;22(2):319-28.
44. Alz.org Facts and Figures. 2015 Alzheimer's Disease Facts and Figures. 2015.
45. Hardy J. A hundred years of Alzheimer's disease research. *Neuron* 2006 Oct 5;52(1):3-13.
46. Duff K, Hardy J. Alzheimer's disease. Mouse model made. *Nature* 1995 Feb 9;373(6514):476-7.
47. Bravard A, Campalans A, Vacher M, Gouget B, Levalois C, Chevillard S, Radicella JP. Inactivation by oxidation and recruitment into stress granules of hOGG1 but not APE1 in human cells exposed to sub-lethal concentrations of cadmium. *Mutat Res* 2010 Mar 1;685(1-2):61-9.
48. Aguzzi A. Neurodegeneration: Alzheimer's disease under strain. *Nature* 2014 Aug 7;512(7512):32-4.
49. Weiner HL, Selkoe DJ. Inflammation and therapeutic vaccination in CNS diseases. *Nature* 2002 Dec 19-26;420(6917):879-84.
50. Licastro F, Carbone I, Raschi E, Porcellini E. The 21st century epidemic: infections as inductors of neuro-degeneration associated with Alzheimer's Disease. *Immun Ageing* 2014;11(1):22.

51. Lambert JC, Ibrahim-Verbaas CA, Harold D, Naj AC, Sims R, Bellenguez C, DeStafano AL, Bis JC, Beecham GW, Grenier-Boley B, Russo G, Thorton-Wells TA, Jones N, Smith AV, Chouraki V, Thomas C, Ikram MA, Zelenika D, Vardarajan BN, Kamatani Y, Lin CF, Gerrish A, Schmidt H, Kunkle B, Dunstan ML, Ruiz A, Bihoreau MT, Choi SH, Reitz C, Pasquier F, Cruchaga C, Craig D, Amin N, Berr C, Lopez OL, De Jager PL, Deramecourt V, Johnston JA, Evans D, Lovestone S, Letenneur L, Moron FJ, Rubinsztein DC, Eiriksdottir G, Sleegers K, Goate AM, Fievet N, Huentelman MW, Gill M, Brown K, Kamboh MI, Keller L, Barberger-Gateau P, McGuinness B, Larson EB, Green R, Myers AJ, Dufouil C, Todd S, Wallon D, Love S, Rogaeva E, Gallacher J, St George-Hyslop P, Clarimon J, Lleo A, Bayer A, Tsuang DW, Yu L, Tsolaki M, Bossu P, Spalletta G, Proitsi P, Collinge J, Sorbi S, Sanchez-Garcia F, Fox NC, Hardy J, Deniz Naranjo MC, Bosco P, Clarke R, Brayne C, Galimberti D, Mancuso M, Matthews F, Moebus S, Mecocci P, Del Zompo M, Maier W, Hampel H, Pilotto A, Bullido M, Panza F, Caffarra P, Nacmias B, Gilbert JR, Mayhaus M, Lannefelt L, Hakonarson H, Pichler S, Carrasquillo MM, Ingelsson M, Beekly D, Alvarez V, Zou F, Valladares O, Younkin SG, Coto E, Hamilton-Nelson KL, Gu W, Razquin C, Pastor P, Mateo I, Owen MJ, Faber KM, Jonsson PV, Combarros O, O'Donovan MC, Cantwell LB, Soininen H, Blacker D, Mead S, Mosley TH, Jr., Bennett DA, Harris TB, Fratiglioni L, Holmes C, de Bruijn RF, Passmore P, Montine TJ, Bettens K, Rotter JI, Brice A, Morgan K, Foroud TM, Kukull WA, Hannequin D, Powell JF, Nalls MA, Ritchie K, Lunetta KL, Kauwe JS, Boerwinkle E, Riemenschneider M, Boada M, Hiltunen M, Martin ER, Schmidt R, Rujescu D, Wang LS, Dartigues JF, Mayeux R, Tzourio C, Hofman A, Nothen MM, Graff C, Psaty BM, Jones L, Haines JL, Holmans PA, Lathrop M, Pericak-Vance MA, Launer LJ, Farrer LA, van Duijn CM, Van Broeckhoven C, Moskvin V, Seshadri S, Williams J, Schellenberg GD, Amouyel P. Meta-analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's disease. *Nat Genet* 2013 Dec;45(12):1452-8.
52. Anderton BH. Alzheimer's disease. Tangled genes and proteins. *Nature* 1987 Sep 10-16;329(6135):106-7.
53. Anderton BH. Alzheimer's disease. Progress in molecular pathology. *Nature* 1987 Feb 19-25;325(6106):658-9.
54. Sahlholm K, Liao F, Holtzman DM, Xu J, Mach RH. Sigma-2 receptor binding is decreased in female, but not male, APP/PS1 mice. *Biochem Biophys Res Commun* 2015 Mar 18.
55. Blennow K, Mattsson N, Scholl M, Hansson O, Zetterberg H. Amyloid biomarkers in Alzheimer's disease. *Trends Pharmacol Sci* 2015 May;36(5):297-309.
56. Sondag CM, Dhawan G, Combs CK. Beta amyloid oligomers and fibrils stimulate differential activation of primary microglia. *J Neuroinflammation* 2009;6:1.
57. Jimenez S, Baglietto-Vargas D, Caballero C, Moreno-Gonzalez I, Torres M, Sanchez-Varo R, Ruano D, Vizuite M, Gutierrez A, Vitorica J. Inflammatory response in the hippocampus of PS1M146L/APP751SL mouse model of Alzheimer's disease: age-dependent switch in the microglial phenotype from alternative to classic. *J Neurosci* 2008 Nov 5;28(45):11650-61.
58. Floden AM, Combs CK. Beta-amyloid stimulates murine postnatal and adult microglia cultures in a unique manner. *J Neurosci* 2006 Apr 26;26(17):4644-8.

59. Dahlgren KN, Manelli AM, Stine WB, Jr., Baker LK, Krafft GA, LaDu MJ. Oligomeric and fibrillar species of amyloid-beta peptides differentially affect neuronal viability. *J Biol Chem* 2002 Aug 30;277(35):32046-53.
60. Selkoe DJ. The molecular pathology of Alzheimer's disease. *Neuron* 1991 Apr;6(4):487-98.
61. Savage MJ, Kalinina J, Wolfe A, Tugusheva K, Korn R, Cash-Mason T, Maxwell JW, Hatcher NG, Haugabook SJ, Wu G, Howell BJ, Renger JJ, Shughrue PJ, McCampbell A. A sensitive abeta oligomer assay discriminates Alzheimer's and aged control cerebrospinal fluid. *J Neurosci* 2014 Feb 19;34(8):2884-97.
62. Mesquita SD, Ferreira AC, Falcao AM, Sousa JC, Oliveira TG, Correia-Neves M, Sousa N, Marques F, Palha JA. Lipocalin 2 modulates the cellular response to amyloid beta. *Cell Death Differ* 2014 Oct;21(10):1588-99.
63. Hardy J, Selkoe DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 2002 Jul 19;297(5580):353-6.
64. Turano E, Busetto G, Marconi S, Guzzo F, Farinazzo A, Commisso M, Bistaffa E, Angiari S, Musumeci S, Sotgiu S, Bonetti B. Neurotoxicity and synaptic plasticity impairment of N-acetylglucosamine polymers: implications for Alzheimer's disease. *Neurobiol Aging* 2015 Jan 6.
65. Jiang T, Tan L, Zhu XC, Zhang QQ, Cao L, Tan MS, Gu LZ, Wang HF, Ding ZZ, Zhang YD, Yu JT. Upregulation of TREM2 ameliorates neuropathology and rescues spatial cognitive impairment in a transgenic mouse model of Alzheimer's disease. *Neuropsychopharmacology* 2014 Dec;39(13):2949-62.
66. Ma J, Choi BR, Chung C, Min SS, Jeon WK, Han JS. Chronic brain inflammation causes a reduction in GluN2A and GluN2B subunits of NMDA receptors and an increase in the phosphorylation of mitogen-activated protein kinases in the hippocampus. *Mol Brain* 2014;7:33.
67. Bachstetter AD, Norris CM, Sompol P, Wilcock DM, Goulding D, Neltner JH, St Clair D, Watterson DM, Van Eldik LJ. Early stage drug treatment that normalizes proinflammatory cytokine production attenuates synaptic dysfunction in a mouse model that exhibits age-dependent progression of Alzheimer's disease-related pathology. *J Neurosci* 2012 Jul 25;32(30):10201-10.
68. McGeer PL, McGeer EG. Targeting microglia for the treatment of Alzheimer's disease. *Expert Opin Ther Targets* 2015 Apr;19(4):497-506.
69. Wisniewski T, Goni F. Immunotherapeutic approaches for Alzheimer's disease. *Neuron* 2015 Mar 18;85(6):1162-76.
70. Canobbio I, Guidetti GF, Oliviero B, Manganaro D, Vara D, Torti M, Pula G. Amyloid beta-peptide-dependent activation of human platelets: essential role for Ca²⁺ and ADP in aggregation and thrombus formation. *Biochem J* 2014 Sep 15;462(3):513-23.
71. Hefendehl JK, Wegenast-Braun BM, Liebig C, Eicke D, Milford D, Calhoun ME, Kohsaka S, Eichner M, Jucker M. Long-term in vivo imaging of beta-amyloid plaque appearance and growth in a mouse model of cerebral beta-amyloidosis. *J Neurosci* 2011 Jan 12;31(2):624-9.

72. Harris-White ME, Chu T, Balverde Z, Sigel JJ, Flanders KC, Frautschy SA. Effects of transforming growth factor-beta (isoforms 1-3) on amyloid-beta deposition, inflammation, and cell targeting in organotypic hippocampal slice cultures. *J Neurosci*1998 Dec 15;18(24):10366-74.
73. Bard F, Cannon C, Barbour R, Burke RL, Games D, Grajeda H, Guido T, Hu K, Huang J, Johnson-Wood K, Khan K, Kholodenko D, Lee M, Lieberburg I, Motter R, Nguyen M, Soriano F, Vasquez N, Weiss K, Welch B, Seubert P, Schenk D, Yednock T. Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. *Nat Med*2000 Aug;6(8):916-9.
74. Cunningham C, Hennessy E. Co-morbidity and systemic inflammation as drivers of cognitive decline: new experimental models adopting a broader paradigm in dementia research. *Alzheimers Res Ther*2015;7(1):33.
75. van Eersel J, Stevens CH, Przybyla M, Gladbach A, Stefanoska K, Chan CK, Ong WY, Hodges JR, Sutherland GT, Kril JJ, Abramowski D, Staufenbiel M, Halliday GM, Ittner LM. Early-onset Axonal Pathology in a Novel P301S-Tau Transgenic Mouse Model of Frontotemporal Lobar Degeneration. *Neuropathol Appl Neurobiol*2015 Mar 12.
76. Spilisbury A, Miwa S, Attems J, Saretzki G. The role of telomerase protein TERT in Alzheimer's disease and in tau-related pathology in vitro. *J Neurosci*2015 Jan 28;35(4):1659-74.
77. Ikezu S, Ikezu T. Tau-tubulin kinase. *Front Mol Neurosci*2014;7:33.
78. Serpente M, Bonsi R, Scarpini E, Galimberti D. Innate immune system and inflammation in Alzheimer's disease: from pathogenesis to treatment. *Neuroimmunomodulation*2014;21(2-3):79-87.
79. Birch AM, Katsouri L, Sastre M. Modulation of inflammation in transgenic models of Alzheimer's disease. *J Neuroinflammation*2014;11:25.
80. Lee S, Xu G, Jay TR, Bhatta S, Kim KW, Jung S, Landreth GE, Ransohoff RM, Lamb BT. Opposing effects of membrane-anchored CX3CL1 on amyloid and tau pathologies via the p38 MAPK pathway. *J Neurosci*2014 Sep 10;34(37):12538-46.
81. Cohen TJ, Guo JL, Hurtado DE, Kwong LK, Mills IP, Trojanowski JQ, Lee VM. The acetylation of tau inhibits its function and promotes pathological tau aggregation. *Nat Commun*2011;2:252.
82. Tremblay MA, Acker CM, Davies P. Tau phosphorylated at tyrosine 394 is found in Alzheimer's disease tangles and can be a product of the Abl-related kinase, Arg. *J Alzheimers Dis*2010;19(2):721-33.
83. Iqbal K, del C. Alonso A, Chen S, Chohan MO, El-Akkad E, Gong C-X, Khatoon S, Li B, Liu F, Rahman A, Tanimukai H, Grundke-Iqbal I. Tau pathology in Alzheimer disease and other tauopathies. *Biochimica et Biophysica Acta (BBA) - Molecular Basis of Disease*2005;1739(2-3):198-210.
84. Mandelkow E. Alzheimer's disease. The tangled tale of tau. *Nature*1999 Dec 9;402(6762):588-9.
85. Lu T, Aron L, Zullo J, Pan Y, Kim H, Chen Y, Yang TH, Kim HM, Drake D, Liu XS, Bennett DA, Colaiacovo MP, Yankner BA. REST and stress resistance in ageing and Alzheimer's disease. *Nature*2014 Mar 27;507(7493):448-54.

86. Wu LY, Bao XQ, Pang HY, Sun H, Zhang D. FLZ attenuates learning and memory deficits via suppressing neuroinflammation induced by LPS in mice. *J Asian Nat Prod Res*2015 Mar;17(3):306-17.
87. Kim EA, Han AR, Choi J, Ahn JY, Choi SY, Cho SW. Anti-inflammatory mechanisms of N-adamantyl-4-methylthiazol-2-amine in lipopolysaccharide-stimulated BV-2 microglial cells. *Int Immunopharmacol*2014 Sep;22(1):73-83.
88. Tegenge MA, Rajbhandari L, Shrestha S, Mithal A, Hosmane S, Venkatesan A. Curcumin protects axons from degeneration in the setting of local neuroinflammation. *Exp Neurol*2014 Mar;253:102-10.
89. Heneka MT, Kummer MP, Stutz A, Delekate A, Schwartz S, Vieira-Saecker A, Griep A, Axt D, Remus A, Tzeng TC, Gelpi E, Halle A, Korte M, Latz E, Golenbock DT. NLRP3 is activated in Alzheimer's disease and contributes to pathology in APP/PS1 mice. *Nature*2013 Jan 31;493(7434):674-8.
90. Sun KH, Chang KH, Clawson S, Ghosh S, Mirzaei H, Regnier F, Shah K. Glutathione-S-transferase P1 is a critical regulator of Cdk5 kinase activity. *J Neurochem*2011 Sep;118(5):902-14.
91. Roy A, Jana A, Yatish K, Freidt MB, Fung YK, Martinson JA, Pahan K. Reactive oxygen species up-regulate CD11b in microglia via nitric oxide: Implications for neurodegenerative diseases. *Free Radic Biol Med*2008 Sep 1;45(5):686-99.
92. Correani V, Francesco LD, Cera I, Mignogna G, Giorgi A, Mazzanti M, Fumagalli L, Fabrizi C, Maras B, Schinina ME. Reversible redox modifications in the microglial proteome challenged by beta amyloid. *Mol Biosyst*2015 Mar 2.
93. Perry G, Cash AD, Smith MA. Alzheimer Disease and Oxidative Stress. *J Biomed Biotechnol*2002;2(3):120-3.
94. Jayasena T, Poljak A, Braidly N, Smythe G, Raftery M, Hill M, Brodaty H, Trollor J, Kochan N, Sachdev P. Upregulation of glycolytic enzymes, mitochondrial dysfunction and increased cytotoxicity in glial cells treated with Alzheimer's disease plasma. *PLoS One*2015;10(3):e0116092.
95. Kuhla A, Ludwig SC, Kuhla B, Munch G, Vollmar B. Advanced glycation end products are mitogenic signals and trigger cell cycle reentry of neurons in Alzheimer's disease brain. *Neurobiol Aging*2015 Feb;36(2):753-61.
96. Swomley AM, Butterfield DA. Oxidative stress in Alzheimer disease and mild cognitive impairment: evidence from human data provided by redox proteomics. *Arch Toxicol*2015 Jul 1.
97. Zetterberg H, Lautner R, Skillback T, Rosen C, Shahim P, Mattsson N, Blennow K. CSF in Alzheimer's disease. *Adv Clin Chem*2014;65:143-72.
98. Urrutia PJ, Mena NP, Nunez MT. The interplay between iron accumulation, mitochondrial dysfunction, and inflammation during the execution step of neurodegenerative disorders. *Front Pharmacol*2014;5:38.
99. Dolzhanskaya N, Merz G, Denman RB. Oxidative stress reveals heterogeneity of FMRP granules in PC12 cell neurites. *Brain Res*2006 Sep 27;1112(1):56-64.
100. Lim GP, Chu T, Yang F, Beech W, Frautschy SA, Cole GM. The curry spice curcumin reduces oxidative damage and amyloid pathology in an Alzheimer transgenic mouse. *J Neurosci*2001 Nov 1;21(21):8370-7.

101. Thinakaran G, Koo EH. Amyloid precursor protein trafficking, processing, and function. *J Biol Chem* 2008 Oct 31;283(44):29615-9.
102. Vassar R. BACE1: the beta-secretase enzyme in Alzheimer's disease. *J Mol Neurosci* 2004;23(1-2):105-14.
103. Pietrzik CU, Yoon IS, Jaeger S, Busse T, Weggen S, Koo EH. FE65 constitutes the functional link between the low-density lipoprotein receptor-related protein and the amyloid precursor protein. *J Neurosci* 2004 Apr 28;24(17):4259-65.
104. Yoshikai S, Sasaki H, Doh-ura K, Furuya H, Sakaki Y. Genomic organization of the human amyloid beta-protein precursor gene. *Gene* 1990 Mar 15;87(2):257-63.
105. Sisodia SS. Beta-amyloid precursor protein cleavage by a membrane-bound protease. *Proc Natl Acad Sci U S A* 1992 Jul 1;89(13):6075-9.
106. Guerreiro RJ, Gustafson DR, Hardy J. The genetic architecture of Alzheimer's disease: beyond APP, PSENs and APOE. *Neurobiol Aging* 2012 Mar;33(3):437-56.
107. Mullan M, Crawford F, Axelman K, Houlden H, Lilius L, Winblad B, Lannfelt L. A pathogenic mutation for probable Alzheimer's disease in the APP gene at the N-terminus of beta-amyloid. *Nat Genet* 1992 Aug;1(5):345-7.
108. Dunn HC, Ager RR, Baglietto-Vargas D, Cheng D, Kitazawa M, Cribbs DH, Medeiros R. Restoration of lipoxin A4 signaling reduces Alzheimer's disease-like pathology in the 3xTg-AD mouse model. *J Alzheimers Dis* 2015;43(3):893-903.
109. De Strooper B. Aph-1, Pen-2, and Nicastrin with Presenilin generate an active gamma-Secretase complex. *Neuron* 2003 Apr 10;38(1):9-12.
110. Rogaev EI, Sherrington R, Rogaeva EA, Levesque G, Ikeda M, Liang Y, Chi H, Lin C, Holman K, Tsuda T, et al. Familial Alzheimer's disease in kindreds with missense mutations in a gene on chromosome 1 related to the Alzheimer's disease type 3 gene. *Nature* 1995 Aug 31;376(6543):775-8.
111. Henricson A, Kall L, Sonnhämmer EL. A novel transmembrane topology of presenilin based on reconciling experimental and computational evidence. *FEBS J* 2005 Jun;272(11):2727-33.
112. Laudon H, Hansson EM, Melen K, Bergman A, Farmery MR, Winblad B, Lendahl U, von Heijne G, Naslund J. A nine-transmembrane domain topology for presenilin 1. *J Biol Chem* 2005 Oct 21;280(42):35352-60.
113. Prox J, Bernreuther C, Altmeyden H, Grendel J, Glatzel M, D'Hooge R, Stroobants S, Ahmed T, Balschun D, Willem M, Lammich S, Isbrandt D, Schweizer M, Horre K, De Strooper B, Saftig P. Postnatal disruption of the disintegrin/metalloproteinase ADAM10 in brain causes epileptic seizures, learning deficits, altered spine morphology, and defective synaptic functions. *J Neurosci* 2013 Aug 7;33(32):12915-28, 28a.
114. Manzine PR, Marcello E, Borroni B, Kamphuis W, Hol E, Padovani A, Nascimento CC, de Godoy Bueno P, Assis Carvalho Vale F, Iost Pavarini SC, Di Luca M, Cominetti MR. ADAM10 gene expression in the blood cells of Alzheimer's disease patients and mild cognitive impairment subjects. *Biomarkers* 2015 Jul 29:1-6.
115. Zeng F, Shen C, Liu YH, Li J, Zhu J, Wang YR, Yan JC, Gao CY, Zhou HD, Deng J, Wang YJ. Genetic association between APP, ADAM10 gene polymorphism, and sporadic Alzheimer's disease in the Chinese population. *Neurotox Res* 2015 Apr;27(3):284-91.

116. Bekris LM, Lutz F, Li G, Galasko DR, Farlow MR, Quinn JF, Kaye JA, Leverenz JB, Tsuang DW, Montine TJ, Peskind ER, Yu CE. ADAM10 expression and promoter haplotype in Alzheimer's disease. *Neurobiol Aging* 2012 Sep;33(9):2229 e1 - e9.
117. Paquet C, Amin J, Mouton-Liger F, Nasser M, Love S, Gray F, Pickering RM, Nicoll JA, Holmes C, Hugon J, Boche D. Effect of active Abeta immunotherapy on neurons in human Alzheimer's disease. *J Pathol* 2015 Apr;235(5):721-30.
118. Chakrabarty P, Li A, Ceballos-Diaz C, Eddy JA, Funk CC, Moore B, DiNunno N, Rosario AM, Cruz PE, Verbeeck C, Sacino A, Nix S, Janus C, Price ND, Das P, Golde TE. IL-10 alters immunoproteostasis in APP mice, increasing plaque burden and worsening cognitive behavior. *Neuron* 2015 Feb 4;85(3):519-33.
119. Rodriguez GA, Tai LM, LaDu MJ, Rebeck GW. Human APOE4 increases microglia reactivity at Abeta plaques in a mouse model of Abeta deposition. *J Neuroinflammation* 2014;11:111.
120. Sriramoju B, Kanwar RK, Kanwar JR. Nanoformulated Mutant SurR9-C84A: a Possible Key for Alzheimer's and its Associated Inflammation. *Pharm Res* 2015 Mar 14.
121. Hu NW, Nicoll AJ, Zhang D, Mably AJ, O'Malley T, Purro SA, Terry C, Collinge J, Walsh DM, Rowan MJ. mGlu5 receptors and cellular prion protein mediate amyloid-beta-facilitated synaptic long-term depression in vivo. *Nat Commun* 2014;5:3374.
122. Richard KL, Filali M, Prefontaine P, Rivest S. Toll-like receptor 2 acts as a natural innate immune receptor to clear amyloid beta 1-42 and delay the cognitive decline in a mouse model of Alzheimer's disease. *J Neurosci* 2008 May 28;28(22):5784-93.
123. Floden AM, Li S, Combs CK. Beta-amyloid-stimulated microglia induce neuron death via synergistic stimulation of tumor necrosis factor alpha and NMDA receptors. *J Neurosci* 2005 Mar 9;25(10):2566-75.
124. Jung CK, Keppler K, Steinbach S, Blazquez-Llorca L, Herms J. Fibrillar amyloid plaque formation precedes microglial activation. *PLoS One* 2015;10(3):e0119768.
125. Ingham V, Williams A, Bate C. Glimepiride reduces CD14 expression and cytokine secretion from macrophages. *J Neuroinflammation* 2014;11:115.
126. Nussbaum JM, Schilling S, Cynis H, Silva A, Swanson E, Wangsanut T, Tayler K, Wiltgen B, Hatami A, Ronicke R, Reymann K, Hutter-Paier B, Alexandru A, Jagla W, Graubner S, Glabe CG, Demuth HU, Bloom GS. Prion-like behaviour and tau-dependent cytotoxicity of pyroglutamylated amyloid-beta. *Nature* 2012 May 31;485(7400):651-5.
127. Glabe C. Intracellular mechanisms of amyloid accumulation and pathogenesis in Alzheimer's disease. *J Mol Neurosci* 2001 Oct;17(2):137-45.
128. Wu Z, Guo Z, Gearing M, Chen G. Tonic inhibition in dentate gyrus impairs long-term potentiation and memory in an Alzheimer's disease model. *Nat Commun* 2014;5:4159.
129. Heneka MT, Ramanathan M, Jacobs AH, Dumitrescu-Ozimek L, Bilkei-Gorzo A, Debeir T, Sastre M, Galldiks N, Zimmer A, Hoehn M, Heiss WD, Klockgether T, Staufenbiel M. Locus ceruleus degeneration promotes Alzheimer pathogenesis in amyloid precursor protein 23 transgenic mice. *J Neurosci* 2006 Feb 1;26(5):1343-54.
130. Heneka MT, Galea E, Gavriluyk V, Dumitrescu-Ozimek L, Daeschner J, O'Banion MK, Weinberg G, Klockgether T, Feinstein DL. Noradrenergic depletion potentiates beta -amyloid-induced cortical inflammation: implications for Alzheimer's disease. *J Neurosci* 2002 Apr 1;22(7):2434-42.

131. Rojanathammanee L, Floden AM, Manocha GD, Combs CK. Attenuation of microglial activation in a mouse model of Alzheimer's disease via NFAT inhibition. *J Neuroinflammation* 2015 Dec;12(1):255.
132. Esteras N, Alquezar C, de la Encarnacion A, Villarejo A, Bermejo-Pareja F, Martin-Requero A. Calmodulin levels in blood cells as a potential biomarker of Alzheimer's disease. *Alzheimers Res Ther* 2013;5(6):55.
133. Min D, Guo F, Zhu S, Xu X, Mao X, Cao Y, Lv X, Gao Q, Wang L, Chen T, Shaw C, Hao L, Cai J. The alterations of Ca²⁺/calmodulin/CaMKII/CaV1.2 signaling in experimental models of Alzheimer's disease and vascular dementia. *Neurosci Lett* 2013 Mar 22;538:60-5.
134. Reese LC, Laezza F, Woltjer R, Taglialatela G. Dysregulated phosphorylation of Ca(2+) /calmodulin-dependent protein kinase II-alpha in the hippocampus of subjects with mild cognitive impairment and Alzheimer's disease. *J Neurochem* 2011 Nov;119(4):791-804.
135. Yoshimura Y, Ichinose T, Yamauchi T. Phosphorylation of tau protein to sites found in Alzheimer's disease brain is catalyzed by Ca²⁺/calmodulin-dependent protein kinase II as demonstrated tandem mass spectrometry. *Neurosci Lett* 2003 Dec 26;353(3):185-8.
136. Combs CK, Johnson DE, Cannady SB, Lehman TM, Landreth GE. Identification of microglial signal transduction pathways mediating a neurotoxic response to amyloidogenic fragments of beta-amyloid and prion proteins. *J Neurosci* 1999 Feb 1;19(3):928-39.
137. Herbst S, Shah A, Mazon Moya M, Marzola V, Jensen B, Reed A, Birrell MA, Saijo S, Mostowy S, Shaunak S, Armstrong-James D. Phagocytosis-dependent activation of a TLR9-BTK-calcineurin-NFAT pathway co-ordinates innate immunity to *Aspergillus fumigatus*. *EMBO Mol Med* 2015;7(3):240-58.
138. Goodridge HS, Simmons RM, Underhill DM. Dectin-1 stimulation by *Candida albicans* yeast or zymosan triggers NFAT activation in macrophages and dendritic cells. *J Immunol* 2007 Mar 1;178(5):3107-15.
139. Wang ZG, Yang C, Zhu B, Hua F. AMPK-dependent autophagic activation is probably involved in the mechanism of resveratrol exerting therapeutic effects for Alzheimer's disease. *Rejuvenation Res* 2015 Feb;18(1):101-2.
140. Vingtdeux V, Davies P, Dickson DW, Marambaud P. AMPK is abnormally activated in tangle- and pre-tangle-bearing neurons in Alzheimer's disease and other tauopathies. *Acta Neuropathol* 2011 Mar;121(3):337-49.
141. Heneka MT, Carson MJ, Khoury JE, Landreth GE, Brosseron F, Feinstein DL, Jacobs AH, Wyss-Coray T, Vitorica J, Ransohoff RM, Herrup K, Frautschy SA, Finsen B, Brown GC, Verkhatsky A, Yamanaka K, Koistinaho J, Latz E, Halle A, Petzold GC, Town T, Morgan D, Shinohara ML, Perry VH, Holmes C, Bazan NG, Brooks DJ, Hunot S, Joseph B, Deigendesch N, Garaschuk O, Boddeke E, Dinarello CA, Breitner JC, Cole GM, Golenbock DT, Kummer MP. Neuroinflammation in Alzheimer's disease. *Lancet Neurol* 2015 Apr;14(4):388-405.
142. Kim EK, Choi EJ. Compromised MAPK signaling in human diseases: an update. *Arch Toxicol* 2015 Feb 18.

143. Gurney ME, D'Amato EC, Burgin AB. Phosphodiesterase-4 (PDE4) molecular pharmacology and Alzheimer's disease. *Neurotherapeutics* 2015 Jan;12(1):49-56.
144. Brawek B, Garaschuk O. Network-wide dysregulation of calcium homeostasis in Alzheimer's disease. *Cell Tissue Res* 2014 Aug;357(2):427-38.
145. Vilchez D, Saez I, Dillin A. The role of protein clearance mechanisms in organismal ageing and age-related diseases. *Nat Commun* 2014;5:5659.
146. Tan L, Yu JT, Tan MS, Liu QY, Wang HF, Zhang W, Jiang T. Genome-wide serum microRNA expression profiling identifies serum biomarkers for Alzheimer's disease. *J Alzheimers Dis* 2014;40(4):1017-27.
147. Funamoto S, Sasaki T, Ishihara S, Nobuhara M, Nakano M, Watanabe-Takahashi M, Saito T, Kakuda N, Miyasaka T, Nishikawa K, Saido TC, Ihara Y. Substrate ectodomain is critical for substrate preference and inhibition of gamma-secretase. *Nat Commun* 2013;4:2529.
148. Frenkel D, Wilkinson K, Zhao L, Hickman SE, Means TK, Puckett L, Farfara D, Kingery ND, Weiner HL, El Khoury J. Scar1 deficiency impairs clearance of soluble amyloid-beta by mononuclear phagocytes and accelerates Alzheimer's-like disease progression. *Nat Commun* 2013;4:2030.
149. Shibuya Y, Chang CC, Huang LH, Bryleva EY, Chang TY. Inhibiting ACAT1/SOAT1 in microglia stimulates autophagy-mediated lysosomal proteolysis and increases Abeta1-42 clearance. *J Neurosci* 2014 Oct 22;34(43):14484-501.
150. Godoy JA, Rios JA, Zolezzi JM, Braidys N, Inestrosa NC. Signaling pathway cross talk in Alzheimer's disease. *Cell Commun Signal* 2014;12:23.
151. Origlia N, Criscuolo C, Arancio O, Yan SS, Domenici L. RAGE inhibition in microglia prevents ischemia-dependent synaptic dysfunction in an amyloid-enriched environment. *J Neurosci* 2014 Jun 25;34(26):8749-60.
152. Chang KH, Multani PS, Sun KH, Vincent F, de Pablo Y, Ghosh S, Gupta R, Lee HP, Lee HG, Smith MA, Shah K. Nuclear envelope dispersion triggered by deregulated Cdk5 precedes neuronal death. *Mol Biol Cell* 2011 May;22(9):1452-62.
153. Chuang JY, Huang YL, Yen WL, Chiang IP, Tsai MH, Tang CH. Syk/JNK/AP-1 signaling pathway mediates interleukin-6-promoted cell migration in oral squamous cell carcinoma. *Int J Mol Sci* 2014;15(1):545-59.
154. Meyerowitz J, Parker SJ, Vella LJ, Ng D, Price KA, Liddell JR, Caragounis A, Li QX, Masters CL, Nonaka T, Hasegawa M, Bogoyevitch MA, Kanninen KM, Crouch PJ, White AR. C-Jun N-terminal kinase controls TDP-43 accumulation in stress granules induced by oxidative stress. *Mol Neurodegener* 2011;6:57.
155. Ploia C, Antoniou X, Scip A, Grande V, Cardinetti D, Colombo A, Canu N, Benussi L, Ghidoni R, Forloni G, Borsello T. JNK plays a key role in tau hyperphosphorylation in Alzheimer's disease models. *J Alzheimers Dis* 2011;26(2):315-29.
156. Mehan S, Meena H, Sharma D, Sankhla R. JNK: a stress-activated protein kinase therapeutic strategies and involvement in Alzheimer's and various neurodegenerative abnormalities. *J Mol Neurosci* 2011 Mar;43(3):376-90.
157. Colombo A, Bastone A, Ploia C, Scip A, Salmona M, Forloni G, Borsello T. JNK regulates APP cleavage and degradation in a model of Alzheimer's disease. *Neurobiol Dis* 2009 Mar;33(3):518-25.

158. Okazawa H, Estus S. The JNK/c-Jun cascade and Alzheimer's disease. *Am J Alzheimers Dis Other Demen*2002 Mar-Apr;17(2):79-88.
159. Scheinfeld MH, Roncarati R, Vito P, Lopez PA, Abdallah M, D'Adamio L. Jun NH2-terminal kinase (JNK) interacting protein 1 (JIP1) binds the cytoplasmic domain of the Alzheimer's beta-amyloid precursor protein (APP). *J Biol Chem*2002 Feb 1;277(5):3767-75.
160. Matsuda S, Yasukawa T, Homma Y, Ito Y, Niikura T, Hiraki T, Hirai S, Ohno S, Kita Y, Kawasumi M, Kouyama K, Yamamoto T, Kyriakis JM, Nishimoto I. c-Jun N-terminal kinase (JNK)-interacting protein-1b/islet-brain-1 scaffolds Alzheimer's amyloid precursor protein with JNK. *J Neurosci*2001 Sep 1;21(17):6597-607.
161. Lopes JP, Oliveira CR, Agostinho P. Neurodegeneration in an Abeta-induced model of Alzheimer's disease: the role of Cdk5. *Aging Cell*2010 Feb;9(1):64-77.
162. Engmann O, Giese KP. Crosstalk between Cdk5 and GSK3beta: Implications for Alzheimer's Disease. *Front Mol Neurosci*2009;2:2.
163. Tsai LH, Lee MS, Cruz J. Cdk5, a therapeutic target for Alzheimer's disease? *Biochim Biophys Acta*2004 Mar 11;1697(1-2):137-42.
164. Cruz JC, Tsai LH. Cdk5 deregulation in the pathogenesis of Alzheimer's disease. *Trends Mol Med*2004 Sep;10(9):452-8.
165. Tandon A, Yu H, Wang L, Rogaeva E, Sato C, Chishti MA, Kawarai T, Hasegawa H, Chen F, Davies P, Fraser PE, Westaway D, St George-Hyslop PH. Brain levels of CDK5 activator p25 are not increased in Alzheimer's or other neurodegenerative diseases with neurofibrillary tangles. *J Neurochem*2003 Aug;86(3):572-81.
166. Pei JJ, Grundke-Iqbal I, Iqbal K, Bogdanovic N, Winblad B, Cowburn RF. Accumulation of cyclin-dependent kinase 5 (cdk5) in neurons with early stages of Alzheimer's disease neurofibrillary degeneration. *Brain Res*1998 Jun 29;797(2):267-77.
167. Jaworski T, Dewachter I, Lechat B, Gees M, Kremer A, Demedts D, Borghgraef P, Devijver H, Kugler S, Patel S, Woodgett JR, Van Leuven F. GSK-3alpha/beta kinases and amyloid production in vivo. *Nature*2011 Dec 8;480(7376):E4-5; discussion E6.
168. Hooper C, Killick R, Lovestone S. The GSK3 hypothesis of Alzheimer's disease. *J Neurochem*2008 Mar;104(6):1433-9.
169. Babcock AA, Ilkjaer L, Clausen BH, Villadsen B, Dissing-Olesen L, Bendixen AT, Lyck L, Lambertsen KL, Finsen B. Cytokine-producing microglia have an altered beta-amyloid load in aged APP/PS1 Tg mice. *Brain Behav Immun*2015 Mar 12.
170. Ward RJ, Dexter DT, Crichton RR. Neurodegenerative diseases and therapeutic strategies using iron chelators. *J Trace Elem Med Biol*2015 Jan 22.
171. Pasqualetti G, Brooks DJ, Edison P. The role of neuroinflammation in dementias. *Curr Neurol Neurosci Rep*2015 Apr;15(4):531.
172. Guillot-Sestier MV, Doty KR, Gate D, Rodriguez J, Jr., Leung BP, Rezai-Zadeh K, Town T. Il10 deficiency rebalances innate immunity to mitigate Alzheimer-like pathology. *Neuron*2015 Feb 4;85(3):534-48.
173. Najjar S, Pearlman DM. Neuroinflammation and white matter pathology in schizophrenia: systematic review. *Schizophr Res*2015 Jan;161(1):102-12.
174. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, Jacobs AH, Wyss-Coray T, Vitorica J, Ransohoff RM, Herrup K, Frautschy SA, Finsen B, Brown GC, Verkhratsky A, Yamanaka K, Koistinaho J, Latz E, Halle A, Petzold GC,

- Town T, Morgan D, Shinohara ML, Perry VH, Holmes C, Bazan NG, Brooks DJ, Hunot S, Joseph B, Deigendesch N, Garaschuk O, Boddeke E, Dinarello CA, Breitner JC, Cole GM, Golenbock DT, Kummer MP. Neuroinflammation in Alzheimer's disease. *Lancet Neurol* 2015 Apr;14(4):388-405.
175. Zhang F, Jiang L. Neuroinflammation in Alzheimer's disease. *Neuropsychiatr Dis Treat* 2015;11:243-56.
176. Malm TM, Jay TR, Landreth GE. The evolving biology of microglia in Alzheimer's disease. *Neurotherapeutics* 2015 Jan;12(1):81-93.
177. Stoeck K, Schmitz M, Ebert E, Schmidt C, Zerr I. Immune responses in rapidly progressive dementia: a comparative study of neuroinflammatory markers in Creutzfeldt-Jakob disease, Alzheimer's disease and multiple sclerosis. *J Neuroinflammation* 2014;11(1):170.
178. Gonzalez H, Elgueta D, Montoya A, Pacheco R. Neuroimmune regulation of microglial activity involved in neuroinflammation and neurodegenerative diseases. *J Neuroimmunol* 2014 Sep 15;274(1-2):1-13.
179. Chen Z, Trapp BD. Microglia and neuroprotection. *J Neurochem* 2015 Feb 18.
180. Cudaback E, Yang Y, Montine TJ, Keene CD. APOE genotype-dependent modulation of astrocyte chemokine CCL3 production. *Glia* 2015 Jan;63(1):51-65.
181. Li Y, Tan MS, Jiang T, Tan L. Microglia in Alzheimer's disease. *Biomed Res Int* 2014;2014:437483.
182. Xu Z, Wu J, Zheng J, Ma H, Zhang H, Zhen X, Zheng LT, Zhang X. Design, synthesis and evaluation of a series of non-steroidal anti-inflammatory drug conjugates as novel neuroinflammatory inhibitors. *Int Immunopharmacol* 2015 Apr;25(2):528-37.
183. Yan Q, Zhang J, Liu H, Babu-Khan S, Vassar R, Biere AL, Citron M, Landreth G. Anti-inflammatory drug therapy alters beta-amyloid processing and deposition in an animal model of Alzheimer's disease. *J Neurosci* 2003 Aug 20;23(20):7504-9.
184. Lim GP, Yang F, Chu T, Chen P, Beech W, Teter B, Tran T, Ubeda O, Ashe KH, Frautschy SA, Cole GM. Ibuprofen suppresses plaque pathology and inflammation in a mouse model for Alzheimer's disease. *J Neurosci* 2000 Aug 1;20(15):5709-14.
185. Suram S, Brown GD, Ghosh M, Gordon S, Loper R, Taylor PR, Akira S, Uematsu S, Williams DL, Leslie CC. Regulation of cytosolic phospholipase A2 activation and cyclooxygenase 2 expression in macrophages by the beta-glucan receptor. *J Biol Chem* 2006 Mar 3;281(9):5506-14.
186. Fujimi K, Noda K, Sasaki K, Wakisaka Y, Tanizaki Y, Iida M, Kiyohara Y, Kanba S, Iwaki T. Altered expression of COX-2 in subdivisions of the hippocampus during aging and in Alzheimer's disease: the Hisayama Study. *Dement Geriatr Cogn Disord* 2007;23(6):423-31.
187. Hoozemans JJ, van Haastert ES, Veerhuis R, Arendt T, Scheper W, Eikelenboom P, Rozemuller AJ. Maximal COX-2 and ppRb expression in neurons occurs during early Braak stages prior to the maximal activation of astrocytes and microglia in Alzheimer's disease. *J Neuroinflammation* 2005 Nov 21;2:27.
188. Xiang Z, Ho L, Valdellon J, Borchelt D, Kelley K, Spielman L, Aisen PS, Pasinetti GM. Cyclooxygenase (COX)-2 and cell cycle activity in a transgenic mouse model of Alzheimer's disease neuropathology. *Neurobiol Aging* 2002 May-Jun;23(3):327-34.

189. Schmitz K, de Bruin N, Bishay P, Mannich J, Haussler A, Altmann C, Ferreiros N, Lotsch J, Ultsch A, Parnham MJ, Geisslinger G, Tegeder I. R-flurbiprofen attenuates experimental autoimmune encephalomyelitis in mice. *EMBO Mol Med* 2014 Nov;6(11):1398-422.
190. Jantzen PT, Connor KE, DiCarlo G, Wenk GL, Wallace JL, Rojiani AM, Coppola D, Morgan D, Gordon MN. Microglial activation and beta -amyloid deposit reduction caused by a nitric oxide-releasing nonsteroidal anti-inflammatory drug in amyloid precursor protein plus presenilin-1 transgenic mice. *J Neurosci* 2002 Mar 15;22(6):2246-54.
191. Choo XY, Alukaidey L, White AR, Grubman A. Neuroinflammation and copper in Alzheimer's disease. *Int J Alzheimers Dis* 2013;2013:145345.
192. Kamphuis W, Kooijman L, Orre M, Stassen O, Pekny M, Hol EM. GFAP and vimentin deficiency alters gene expression in astrocytes and microglia in wild-type mice and changes the transcriptional response of reactive glia in mouse model for Alzheimer's disease. *Glia* 2015 Mar 2.
193. Bataveljic D, Petrovic J, Lazic K, Saponjic J, Andjus P. Glial response in the rat models of functionally distinct cholinergic neuronal denervations. *J Neurosci Res* 2015 Feb;93(2):244-52.
194. Oberstein TJ, Spitzer P, Klafki HW, Linning P, Neff F, Knolker HJ, Lewczuk P, Wiltfang J, Kornhuber J, Maler JM. Astrocytes and microglia but not neurons preferentially generate N-terminally truncated Abeta peptides. *Neurobiol Dis* 2015 Jan;73:24-35.
195. Phillips EC, Croft CL, Kurbatskaya K, O'Neill MJ, Hutton ML, Hanger DP, Garwood CJ, Noble W. Astrocytes and neuroinflammation in Alzheimer's disease. *Biochem Soc Trans* 2014 Oct;42(5):1321-5.
196. Hancock SM, Finkelstein DI, Adlard PA. Glia and zinc in ageing and Alzheimer's disease: a mechanism for cognitive decline? *Front Aging Neurosci* 2014;6:137.
197. Jung ES, An K, Hong HS, Kim JH, Mook-Jung I. Astrocyte-originated ATP protects Abeta(1-42)-induced impairment of synaptic plasticity. *J Neurosci* 2012 Feb 29;32(9):3081-7.
198. Gao XF, Wang W, Yu Q, Burnstock G, Xiang ZH, He C. Astroglial P2X7 receptor current density increased following long-term exposure to rotenone. *Purinergic Signal* 2011 Mar;7(1):65-72.
199. Dickson DW, Mattiace LA. Astrocytes and microglia in human brain share an epitope recognized by a B-lymphocyte-specific monoclonal antibody (LN-1). *Am J Pathol* 1989 Jul;135(1):135-47.
200. Filipcik P, Cente M, Zilka N, Smolek T, Hanes J, Kucerak J, Opattova A, Kovacech B, Novak M. Intraneuronal accumulation of misfolded tau protein induces overexpression of Hsp27 in activated astrocytes. *Biochim Biophys Acta* 2015 Mar 12.
201. Ko SY, Lee HE, Park SJ, Jeon SJ, Kim B, Gao Q, Jang DS, Ryu JH. Spinosin, a C-Glucosylflavone, from *Zizyphus jujuba* var. *spinosa* Ameliorates Abeta1-42 Oligomer-Induced Memory Impairment in Mice. *Biomol Ther (Seoul)* 2015 Mar;23(2):156-64.
202. Lee M, Shi X, Barron AE, McGeer E, McGeer PL. Human antimicrobial peptide LL-37 induces glial-mediated neuroinflammation. *Biochem Pharmacol* 2015 Mar 15;94(2):130-41.

203. Wilcock DM, Vitek MP, Colton CA. Vascular amyloid alters astrocytic water and potassium channels in mouse models and humans with Alzheimer's disease. *Neuroscience* 2009 Mar 31;159(3):1055-69.
204. Luna-Medina R, Cortes-Canteli M, Sanchez-Galiano S, Morales-Garcia JA, Martinez A, Santos A, Perez-Castillo A. NP031112, a thiadiazolidinone compound, prevents inflammation and neurodegeneration under excitotoxic conditions: potential therapeutic role in brain disorders. *J Neurosci* 2007 May 23;27(21):5766-76.
205. Jacobsen JS, Wu CC, Redwine JM, Comery TA, Arias R, Bowlby M, Martone R, Morrison JH, Pangalos MN, Reinhart PH, Bloom FE. Early-onset behavioral and synaptic deficits in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A* 2006 Mar 28;103(13):5161-6.
206. Phillips HS, Hains JM, Armanini M, Laramie GR, Johnson SA, Winslow JW. BDNF mRNA is decreased in the hippocampus of individuals with Alzheimer's disease. *Neuron* 1991 Nov;7(5):695-702.
207. Harvey H, Durant S. The role of glial cells and the complement system in retinal diseases and Alzheimer's disease: common neural degeneration mechanisms. *Exp Brain Res* 2014 Nov;232(11):3363-77.
208. Nguyen TV, Shen L, Vander Griend L, Quach LN, Belichenko NP, Saw N, Yang T, Shamloo M, Wyss-Coray T, Massa SM, Longo FM. Small molecule p75NTR ligands reduce pathological phosphorylation and misfolding of tau, inflammatory changes, cholinergic degeneration, and cognitive deficits in AbetaPP(L/S) transgenic mice. *J Alzheimers Dis* 2014;42(2):459-83.
209. Sawyer AJ, Tian W, Saucier-Sawyer JK, Rizk PJ, Saltzman WM, Bellamkonda RV, Kyriakides TR. The effect of inflammatory cell-derived MCP-1 loss on neuronal survival during chronic neuroinflammation. *Biomaterials* 2014 Aug;35(25):6698-706.
210. Ho GJ, Drego R, Hakimian E, Masliah E. Mechanisms of cell signaling and inflammation in Alzheimer's disease. *Curr Drug Targets Inflamm Allergy* 2005 Apr;4(2):247-56.
211. Nayak D, Roth TL, McGavern DB. Microglia development and function. *Annu Rev Immunol* 2014;32:367-402.
212. Barron KD. The microglial cell. A historical review. *J Neurol Sci* 1995 Dec;134 Suppl:57-68.
213. Latta CH, Brothers HM, Wilcock DM. Neuroinflammation in Alzheimer's disease; A source of heterogeneity and target for personalized therapy. *Neuroscience* 2014 Oct 5.
214. Saijo K, Glass CK. Microglial cell origin and phenotypes in health and disease. *Nat Rev Immunol* 2011 Nov;11(11):775-87.
215. Navascues J, Moujahid A, Almendros A, Marin-Teva JL, Cuadros MA. Origin of microglia in the quail retina: central-to-peripheral and vitreal-to-scleral migration of microglial precursors during development. *J Comp Neurol* 1995 Apr 3;354(2):209-28.
216. Ling EA, Wong WC. The origin and nature of ramified and amoeboid microglia: a historical review and current concepts. *Glia* 1993 Jan;7(1):9-18.
217. Macauley SL, Wong AM, Shyng C, Augner DP, Dearborn JT, Pearse Y, Roberts MS, Fowler SC, Cooper JD, Watterson DM, Sands MS. An anti-neuroinflammatory that

- targets dysregulated glia enhances the efficacy of CNS-directed gene therapy in murine infantile neuronal ceroid lipofuscinosis. *J Neurosci*2014 Sep 24;34(39):13077-82.
218. Kudo M. Environmental Pathology: SY09-2 MULTIPLE SCLEROSIS (MS) AND NEURODEGENERATION: CAUSE AND PATHOGENESIS IN RELATION TO ELECTROMAGNETIC FIELDS (EMF). *Pathology*2014 Oct;46 Suppl 2:S15.
219. Streit WJ, Xue QS. Human CNS immune senescence and neurodegeneration. *Curr Opin Immunol*2014 Aug;29:93-6.
220. Doens D, Fernandez PL. Microglia receptors and their implications in the response to amyloid beta for Alzheimer's disease pathogenesis. *J Neuroinflammation*2014;11:48.
221. Harry GJ. Microglia during development and aging. *Pharmacol Ther*2013 Sep;139(3):313-26.
222. Krauthausen M, Kummer MP, Zimmermann J, Reyes-Irisarri E, Terwel D, Bulic B, Heneka MT, Muller M. CXCR3 promotes plaque formation and behavioral deficits in an Alzheimer's disease model. *J Clin Invest*2015 Jan;125(1):365-78.
223. Shah VB, Huang Y, Keshwara R, Ozment-Skelton T, Williams DL, Keshvara L. Beta-glucan activates microglia without inducing cytokine production in Dectin-1-dependent manner. *J Immunol*2008 Mar 1;180(5):2777-85.
224. Song X, Tanaka S, Cox D, Lee SC. Fcgamma receptor signaling in primary human microglia: differential roles of PI-3K and Ras/ERK MAPK pathways in phagocytosis and chemokine induction. *J Leukoc Biol*2004 Jun;75(6):1147-55.
225. Yu Y, Ye RD. Microglial Abeta receptors in Alzheimer's disease. *Cell Mol Neurobiol*2015 Jan;35(1):71-83.
226. Takahashi K. [Neurodegenerative disease and TREM2]. *Rinsho Shinkeigaku*2014;54(12):1122-4.
227. Larochelle A, Bellavance MA, Rivest S. Role of adaptor protein MyD88 in TLR-mediated preconditioning and neuroprotection after acute excitotoxicity. *Brain Behav Immun*2015 Feb 27.
228. Gambuzza ME, Sofo V, Salmeri FM, Soraci L, Marino S, Bramanti P. Toll-like receptors in Alzheimer's disease: a therapeutic perspective. *CNS Neurol Disord Drug Targets*2014;13(9):1542-58.
229. Liu T, Ji RR. Toll-Like Receptors and Itch. 2014.
230. Reed-Geaghan EG, Savage JC, Hise AG, Landreth GE. CD14 and toll-like receptors 2 and 4 are required for fibrillar A{beta}-stimulated microglial activation. *J Neurosci*2009 Sep 23;29(38):11982-92.
231. Ma J, Jiang T, Tan L, Yu JT. TYROBP in Alzheimer's Disease. *Mol Neurobiol*2015 Apr;51(2):820-6.
232. Fuller JP, Stavenhagen JB, Teeling JL. New roles for Fc receptors in neurodegeneration-the impact on Immunotherapy for Alzheimer's Disease. *Front Neurosci*2014;8:235.
233. Santiago AR, Baptista FI, Santos PF, Cristovao G, Ambrosio AF, Cunha RA, Gomes CA. Role of microglia adenosine A(2A) receptors in retinal and brain neurodegenerative diseases. *Mediators Inflamm*2014;2014:465694.
234. Fu R, Shen Q, Xu P, Luo JJ, Tang Y. Phagocytosis of microglia in the central nervous system diseases. *Mol Neurobiol*2014 Jun;49(3):1422-34.

235. Hickman SE, El Khoury J. TREM2 and the neuroimmunology of Alzheimer's disease. *Biochem Pharmacol*2014 Apr 15;88(4):495-8.
236. Lucin KM, O'Brien CE, Bieri G, Czirr E, Mosher KI, Abbey RJ, Mastroeni DF, Rogers J, Spencer B, Masliah E, Wyss-Coray T. Microglial beclin 1 regulates retromer trafficking and phagocytosis and is impaired in Alzheimer's disease. *Neuron*2013 Sep 4;79(5):873-86.
237. Naert G, Rivest S. CC chemokine receptor 2 deficiency aggravates cognitive impairments and amyloid pathology in a transgenic mouse model of Alzheimer's disease. *J Neurosci*2011 Apr 20;31(16):6208-20.
238. Takayama F, Wu Z, Ma HM, Okada R, Hayashi Y, Nakanishi H. Possible involvement of aiPLA2 in the phosphatidylserine-containing liposomes induced production of PGE2 and PGD2 in microglia. *J Neuroimmunol*2013 Sep 15;262(1-2):121-4.
239. Chang GH, Barbaro NM, Pieper RO. Phosphatidylserine-dependent phagocytosis of apoptotic glioma cells by normal human microglia, astrocytes, and glioma cells. *Neuro Oncol*2000 Jul;2(3):174-83.
240. Witting A, Muller P, Herrmann A, Kettenmann H, Nolte C. Phagocytic clearance of apoptotic neurons by Microglia/Brain macrophages in vitro: involvement of lectin-, integrin-, and phosphatidylserine-mediated recognition. *J Neurochem*2000 Sep;75(3):1060-70.
241. Mazaheri F, Breus O, Durdu S, Haas P, Wittbrodt J, Gilmour D, Peri F. Distinct roles for BAI1 and TIM-4 in the engulfment of dying neurons by microglia. *Nat Commun*2014;5:4046.
242. Grizenkova J, Akhtar S, Brandner S, Collinge J, Lloyd SE. Microglial Cx3cr1 knockout reduces prion disease incubation time in mice. *BMC Neurosci*2014;15:44.
243. Burnstock G. Physiopathological roles of P2X receptors in the central nervous system. *Curr Med Chem*2015;22(7):819-44.
244. Delekate A, Fuchtemeier M, Schumacher T, Ulbrich C, Foddiss M, Petzold GC. Metabotropic P2Y1 receptor signalling mediates astrocytic hyperactivity in vivo in an Alzheimer's disease mouse model. *Nat Commun*2014;5:5422.
245. Lue LF, Schmitz C, Walker DG. What happens to microglial TREM2 in Alzheimer's disease: Immunoregulatory turned into immunopathogenic? *Neuroscience*2014 Oct 2.
246. Graham LM, Gupta V, Schafer G, Reid DM, Kimberg M, Dennehy KM, Hornsell WG, Guler R, Campanero-Rhodes MA, Palma AS, Feizi T, Kim SK, Sobieszczuk P, Willment JA, Brown GD. The C-type lectin receptor CLECSF8 (CLEC4D) is expressed by myeloid cells and triggers cellular activation through Syk kinase. *J Biol Chem*2012 Jul 27;287(31):25964-74.
247. Johansson JU, Woodling NS, Wang Q, Panchal M, Liang X, Trueba-Saiz A, Brown HD, Mhatre SD, Loui T, Andreasson KI. Prostaglandin signaling suppresses beneficial microglial function in Alzheimer's disease models. *J Clin Invest*2015 Jan;125(1):350-64.
248. Parajuli B, Sonobe Y, Horiuchi H, Takeuchi H, Mizuno T, Suzumura A. Oligomeric amyloid beta induces IL-1beta processing via production of ROS: implication in Alzheimer's disease. *Cell Death Dis*2013;4:e975.

249. Kim YJ, Hwang SY, Oh ES, Oh S, Han IO. IL-1beta, an immediate early protein secreted by activated microglia, induces iNOS/NO in C6 astrocytoma cells through p38 MAPK and NF-kappaB pathways. *J Neurosci Res* 2006 Oct;84(5):1037-46.
250. Kyrkanides S, Olschowka JA, Williams JP, Hansen JT, O'Banion MK. TNF alpha and IL-1beta mediate intercellular adhesion molecule-1 induction via microglia-astrocyte interaction in CNS radiation injury. *J Neuroimmunol* 1999 Mar 1;95(1-2):95-106.
251. Bhat RV, DiRocco R, Marcy VR, Flood DG, Zhu Y, Dobrzanski P, Siman R, Scott R, Contreras PC, Miller M. Increased expression of IL-1beta converting enzyme in hippocampus after ischemia: selective localization in microglia. *J Neurosci* 1996 Jul 1;16(13):4146-54.
252. Barrett JP, Minogue AM, Jones RS, Ribeiro C, Kelly RJ, Lynch MA. Bone marrow-derived macrophages from AbetaPP/PS1 mice are sensitized to the effects of inflammatory stimuli. *J Alzheimers Dis* 2015 Jan 1;44(3):949-62.
253. Minogue AM, Jones RS, Kelly RJ, McDonald CL, Connor TJ, Lynch MA. Age-associated dysregulation of microglial activation is coupled with enhanced blood-brain barrier permeability and pathology in APP/PS1 mice. *Neurobiol Aging* 2014 Jun;35(6):1442-52.
254. Gaikwad S, Larionov S, Wang Y, Dannenberg H, Matozaki T, Monsonego A, Thal DR, Neumann H. Signal regulatory protein-beta1: a microglial modulator of phagocytosis in Alzheimer's disease. *Am J Pathol* 2009 Dec;175(6):2528-39.
255. Meda L, Cassatella MA, Szendrei GI, Otvos L, Jr., Baron P, Villalba M, Ferrari D, Rossi F. Activation of microglial cells by beta-amyloid protein and interferon-gamma. *Nature* 1995 Apr 13;374(6523):647-50.
256. Duan S, Guan X, Lin R, Liu X, Yan Y, Zhang T, Chen X, Huang J, Sun X, Li Q, Fang S, Xu J, Yao Z, Gu H. Silibinin inhibits acetylcholinesterase activity and amyloid beta peptide aggregation: a dual-target drug for the treatment of Alzheimer's disease. *Neurobiol Aging* 2015 Feb 11.
257. Moon M, Cha MY, Mook-Jung I. Impaired hippocampal neurogenesis and its enhancement with ghrelin in 5XFAD mice. *J Alzheimers Dis* 2014;41(1):233-41.
258. Goldmann T, Tay TL, Prinz M. Love and death: microglia, NLRP3 and the Alzheimer's brain. *Cell Res* 2013 May;23(5):595-6.
259. Scheib JL, Sullivan CS, Carter BD. Jedi-1 and MEGF10 signal engulfment of apoptotic neurons through the tyrosine kinase Syk. *J Neurosci* 2012 Sep 19;32(38):13022-31.
260. Oster-Granite ML, McPhie DL, Greenan J, Neve RL. Age-dependent neuronal and synaptic degeneration in mice transgenic for the C terminus of the amyloid precursor protein. *J Neurosci* 1996 Nov 1;16(21):6732-41.
261. Bahrini I, Song JH, Diez D, Hanayama R. Neuronal exosomes facilitate synaptic pruning by up-regulating complement factors in microglia. *Sci Rep* 2015;5:7989.
262. Paolicelli RC, Bolasco G, Pagani F, Maggi L, Scianni M, Panzanelli P, Giustetto M, Ferreira TA, Guiducci E, Dumas L, Ragozzino D, Gross CT. Synaptic pruning by microglia is necessary for normal brain development. *Science* 2011 Sep 9;333(6048):1456-8.

263. Michailidou I, Willems JG, Kooi EJ, van Eden C, Gold SM, Geurts JJ, Baas F, Huitinga I, Ramaglia V. Complement C1q-C3 associated synaptic changes in multiple sclerosis hippocampus. *Ann Neurol* 2015 Feb 27.
264. Wes PD, Holtman IR, Boddeke EW, Moller T, Eggen BJ. Next generation transcriptomics and genomics elucidate biological complexity of microglia in health and disease. *Glia* 2015 Jun 4.
265. Hadas S, Spira M, Hanisch UK, Reichert F, Rotshenker S. Complement receptor-3 negatively regulates the phagocytosis of degenerated myelin through tyrosine kinase Syk and cofilin. *J Neuroinflammation* 2012;9:166.
266. Maier M, Peng Y, Jiang L, Seabrook TJ, Carroll MC, Lemere CA. Complement C3 deficiency leads to accelerated amyloid beta plaque deposition and neurodegeneration and modulation of the microglia/macrophage phenotype in amyloid precursor protein transgenic mice. *J Neurosci* 2008 Jun 18;28(25):6333-41.
267. Roussos P, Katsel P, Fam P, Tan W, Purohit DP, Haroutunian V. The triggering receptor expressed on myeloid cells 2 (TREM2) is associated with enhanced inflammation, neuropathological lesions and increased risk for Alzheimer's dementia. *Alzheimers Dement* 2014 Dec 9.
268. Kleinberger G, Yamanishi Y, Suarez-Calvet M, Czirr E, Lohmann E, Cuyvers E, Struyfs H, Pettkus N, Wenninger-Weinzierl A, Mazaheri F, Tahirovic S, Lleo A, Alcolea D, Fortea J, Willem M, Lammich S, Molinuevo JL, Sanchez-Valle R, Antonell A, Ramirez A, Heneka MT, Slegers K, van der Zee J, Martin JJ, Engelborghs S, Demirtas-Tatlidede A, Zetterberg H, Van Broeckhoven C, Gurvit H, Wyss-Coray T, Hardy J, Colonna M, Haass C. TREM2 mutations implicated in neurodegeneration impair cell surface transport and phagocytosis. *Sci Transl Med* 2014 Jul 2;6(243):243ra86.
269. Satoh J, Asahina N, Kitano S, Kino Y. A Comprehensive Profile of ChIP-Seq-Based PU.1/Spi1 Target Genes in Microglia. *Gene Regul Syst Bio* 2014;8:127-39.
270. Gomez-Nicola D, Fransen NL, Suzzi S, Perry VH. Regulation of microglial proliferation during chronic neurodegeneration. *J Neurosci* 2013 Feb 6;33(6):2481-93.
271. Yi YS, Son YJ, Ryou C, Sung GH, Kim JH, Cho JY. Functional roles of Syk in macrophage-mediated inflammatory responses. *Mediators Inflamm* 2014;2014:270302.
272. Wang CY, Wang ZY, Xie JW, Cai JH, Wang T, Xu Y, Wang X, An L. CD36 upregulation mediated by intranasal LV-NRF2 treatment mitigates hypoxia-induced progression of Alzheimer's-like pathogenesis. *Antioxid Redox Signal* 2014 Dec 1;21(16):2208-30.
273. Koenigsknecht J, Landreth G. Microglial phagocytosis of fibrillar beta-amyloid through a beta1 integrin-dependent mechanism. *J Neurosci* 2004 Nov 3;24(44):9838-46.
274. Joshi P, Turola E, Ruiz A, Bergami A, Libera DD, Benussi L, Giussani P, Magnani G, Comi G, Legname G, Ghidoni R, Furlan R, Matteoli M, Verderio C. Microglia convert aggregated amyloid-beta into neurotoxic forms through the shedding of microvesicles. *Cell Death Differ* 2014 Apr;21(4):582-93.
275. Hussain SF, Kong LY, Jordan J, Conrad C, Madden T, Fokt I, Priebe W, Heimberger AB. A novel small molecule inhibitor of signal transducers and activators of transcription 3 reverses immune tolerance in malignant glioma patients. *Cancer Res* 2007 Oct 15;67(20):9630-6.

276. Blasko I, Stampfer-Kountchev M, Robatscher P, Veerhuis R, Eikelenboom P, Grubeck-Loebenstien B. How chronic inflammation can affect the brain and support the development of Alzheimer's disease in old age: the role of microglia and astrocytes. *Aging Cell*2004 Aug;3(4):169-76.
277. Fakhoury M. Role of Immunity and Inflammation in the Pathophysiology of Neurodegenerative Diseases. *Neurodegener Dis*2015 Jan 15.
278. Michaud JP, Rivest S. Anti-inflammatory signaling in microglia exacerbates Alzheimer's disease-related pathology. *Neuron*2015 Feb 4;85(3):450-2.
279. Marlatt MW, Bauer J, Aronica E, van Haastert ES, Hoozemans JJ, Joels M, Lucassen PJ. Proliferation in the Alzheimer hippocampus is due to microglia, not astroglia, and occurs at sites of amyloid deposition. *Neural Plast*2014;2014:693851.
280. Zolezzi JM, Bastias-Candia S, Santos MJ, Inestrosa NC. Alzheimer's disease: relevant molecular and physiopathological events affecting amyloid-beta brain balance and the putative role of PPARs. *Front Aging Neurosci*2014;6:176.
281. Portelius E, Soininen H, Andreasson U, Zetterberg H, Persson R, Karlsson G, Blennow K, Herukka SK, Mattsson N. Exploring Alzheimer molecular pathology in Down's syndrome cerebrospinal fluid. *Neurodegener Dis*2014;14(2):98-106.
282. Streit WJ, Xue QS, Tischer J, Bechmann I. Microglial pathology. *Acta Neuropathol Commun*2014;2:142.
283. Streit WJ, Xue QS. Life and death of microglia. *J Neuroimmune Pharmacol*2009 Dec;4(4):371-9.
284. Zhang H, Su YJ, Zhou WW, Wang SW, Xu PX, Yu XL, Liu RT. Activated scavenger receptor A promotes glial internalization of abeta. *PLoS One*2014;9(4):e94197.
285. Mandrekar S, Jiang Q, Lee CY, Koenigsnecht-Talboo J, Holtzman DM, Landreth GE. Microglia mediate the clearance of soluble Abeta through fluid phase macropinocytosis. *J Neurosci*2009 Apr 1;29(13):4252-62.
286. Paradowska-Gorycka A, Jurkowska M. Structure, expression pattern and biological activity of molecular complex TREM-2/DAP12. *Hum Immunol*2013 Jun;74(6):730-7.
287. Satoh J, Tabunoki H, Ishida T, Yagishita S, Jinnai K, Futamura N, Kobayashi M, Toyoshima I, Yoshioka T, Enomoto K, Arai N, Saito Y, Arima K. Phosphorylated Syk expression is enhanced in Nasu-Hakola disease brains. *Neuropathology*2012 Apr;32(2):149-57.
288. Walker DG, Whetzel AM, Serrano G, Sue LI, Beach TG, Lue LF. Association of CD33 polymorphism rs3865444 with Alzheimer's disease pathology and CD33 expression in human cerebral cortex. *Neurobiol Aging*2015 Feb;36(2):571-82.
289. Bradshaw EM, Chibnik LB, Keenan BT, Ottoboni L, Raj T, Tang A, Rosenkrantz LL, Imboywa S, Lee M, Von Korff A, Morris MC, Evans DA, Johnson K, Sperling RA, Schneider JA, Bennett DA, De Jager PL. CD33 Alzheimer's disease locus: altered monocyte function and amyloid biology. *Nat Neurosci*2013 Jul;16(7):848-50.
290. Malik M, Simpson JF, Parikh I, Wilfred BR, Fardo DW, Nelson PT, Estus S. CD33 Alzheimer's risk-altering polymorphism, CD33 expression, and exon 2 splicing. *J Neurosci*2013 Aug 14;33(33):13320-5.

291. Geahlen RL. Getting Syk: spleen tyrosine kinase as a therapeutic target. *Trends Pharmacol Sci*2014 Aug;35(8):414-22.
292. Geahlen RL, Burg DL. The role of Syk in cell signaling. *Adv Exp Med Biol*1994;365:103-9.
293. Brungs S, Kolanus W, Hemmersbach R. Syk phosphorylation inverted question mark a gravesensitive step in macrophage signalling. *Cell Commun Signal*2015 Feb 3;13(1):9.
294. Kimura Y, Chihara K, Honjoh C, Takeuchi K, Yamauchi S, Yoshiki H, Fujieda S, Sada K. Dectin-1-mediated signaling leads to characteristic gene expressions and cytokine secretion via spleen tyrosine kinase (Syk) in rat mast cells. *J Biol Chem*2014 Nov 7;289(45):31565-75.
295. Canetti C, Hu B, Curtis JL, Peters-Golden M. Syk activation is a leukotriene B4-regulated event involved in macrophage phagocytosis of IgG-coated targets but not apoptotic cells. *Blood*2003 Sep 1;102(5):1877-83.
296. Turner M, Mee PJ, Costello PS, Williams O, Price AA, Duddy LP, Furlong MT, Geahlen RL, Tybulewicz VL. Perinatal lethality and blocked B-cell development in mice lacking the tyrosine kinase Syk. *Nature*1995 Nov 16;378(6554):298-302.
297. Tan SL, Liao C, Lucas MC, Stevenson C, DeMartino JA. Targeting the SYK-BTK axis for the treatment of immunological and hematological disorders: recent progress and therapeutic perspectives. *Pharmacol Ther*2013 May;138(2):294-309.
298. Ozaki N, Suzuki S, Ishida M, Harada Y, Tanaka K, Sato Y, Kono T, Kubo M, Kitamura D, Encinas J, Hara H, Yoshida H. Syk-dependent signaling pathways in neutrophils and macrophages are indispensable in the pathogenesis of anti-collagen antibody-induced arthritis. *Int Immunol*2012 Sep;24(9):539-50.
299. Iwata S, Nakayamada S, Fukuyo S, Kubo S, Yunoue N, Wang SP, Yoshikawa M, Saito K, Tanaka Y. Activation of Syk in peripheral blood B cells in patients with rheumatoid arthritis: a potential target for abatacept therapy. *Arthritis Rheumatol*2015 Jan;67(1):63-73.
300. Jiang M, Liu R, Chen Y, Zheng Q, Fan S, Liu P. A combined experimental and computational study of Vam3, a derivative of resveratrol, and Syk interaction. *Int J Mol Sci*2014;15(9):17188-203.
301. Nemeth T, Mocsai A. The role of neutrophils in autoimmune diseases. *Immunol Lett*2012 Mar 30;143(1):9-19.
302. Bajpai M. Fostamatinib, a Syk inhibitor prodrug for the treatment of inflammatory diseases. *IDrugs*2009 Mar;12(3):174-85.
303. Behnen M, Leschczyk C, Moller S, Batel T, Klinger M, Solbach W, Laskay T. Immobilized immune complexes induce neutrophil extracellular trap release by human neutrophil granulocytes via FcγRIIIB and Mac-1. *J Immunol*2014 Aug 15;193(4):1954-65.
304. Gradler U, Schwarz D, Dresing V, Musil D, Bomke J, Frech M, Greiner H, Jakel S, Rysiok T, Muller-Pompalla D, Wegener A. Structural and biophysical characterization of the Syk activation switch. *J Mol Biol*2013 Jan 23;425(2):309-33.
305. Kaur M, Singh M, Silakari O. Inhibitors of switch kinase 'spleen tyrosine kinase' in inflammation and immune-mediated disorders: a review. *Eur J Med Chem*2013 Sep;67:434-46.

306. Barr PM, Wei C, Roger J, Schaefer-Cutillo J, Kelly JL, Rosenberg AF, Jung J, Sanz I, Friedberg JW. Syk inhibition with fostamatinib leads to transitional B lymphocyte depletion. *Clin Immunol* 2012 Mar;142(3):237-42.
307. Hossen MJ, Baek KS, Kim E, Yang WS, Jeong D, Kim JH, Kweon DH, Yoon DH, Kim TW, Cho JY. In vivo and in vitro anti-inflammatory activities of *Persicaria chinensis* methanolic extract targeting Src/Syk/NF-kappaB. *J Ethnopharmacol* 2015 Jan 15;159:9-16.
308. Sithanandam G, Anderson LM. The ERBB3 receptor in cancer and cancer gene therapy. *Cancer Gene Ther* 2008 Jul;15(7):413-48.
309. Parsa KV, Butchar JP, Rajaram MV, Cremer TJ, Tridandapani S. The tyrosine kinase Syk promotes phagocytosis of *Francisella* through the activation of Erk. *Mol Immunol* 2008 May;45(10):3012-21.
310. Khandani A, Eng E, Jongstra-Bilen J, Schreiber AD, Douda D, Samavarchi-Tehrani P, Harrison RE. Microtubules regulate PI-3K activity and recruitment to the phagocytic cup during Fcgamma receptor-mediated phagocytosis in nonelicited macrophages. *J Leukoc Biol* 2007 Aug;82(2):417-28.
311. Paris D, Ait-Ghezala G, Bachmeier C, Laco G, Beaulieu-Abdelahad D, Lin Y, Jin C, Crawford F, Mullan M. The spleen tyrosine kinase (Syk) regulates Alzheimer amyloid-beta production and Tau hyperphosphorylation. *J Biol Chem* 2014 Dec 5;289(49):33927-44.
312. Cui S, Bilitewski U. [Anti-inflammatory effect of Syk inhibitor in LPS stimulated macrophages]. *Xi Bao Yu Fen Zi Mian Yi Xue Za Zhi* 2013 Oct;29(10):1024-7.
313. Gallagher D, Gutierrez H, Gavalda N, O'Keeffe G, Hay R, Davies AM. Nuclear factor-kappaB activation via tyrosine phosphorylation of inhibitor kappaB-alpha is crucial for ciliary neurotrophic factor-promoted neurite growth from developing neurons. *J Neurosci* 2007 Sep 5;27(36):9664-9.
314. Yankee TM, Keshvara LM, Sawasdikosol S, Harrison ML, Geahlen RL. Inhibition of signaling through the B cell antigen receptor by the protooncogene product, c-Cbl, requires Syk tyrosine 317 and the c-Cbl phosphotyrosine-binding domain. *J Immunol* 1999 Dec 1;163(11):5827-35.
315. Manne BK, Badolia R, Dangelmaier C, Eble JA, Ellmeier W, Kahn M, Kunapuli SP. Distinct pathways regulate Syk activation downstream of ITAM and hemITAM receptors in platelets. *J Biol Chem* 2015 Mar 12.
316. Sada K, Takano T, Yanagi S, Yamamura H. Structure and function of Syk protein-tyrosine kinase. *J Biochem* 2001 Aug;130(2):177-86.
317. Krisenko MO, Geahlen RL. Calling in SYK: SYK's dual role as a tumor promoter and tumor suppressor in cancer. *Biochim Biophys Acta* 2015 Jan;1853(1):254-63.
318. Zhao H, Caflisch A. Discovery of dual ZAP70 and Syk kinases inhibitors by docking into a rare C-helix-out conformation of Syk. *Bioorg Med Chem Lett* 2014 Mar 15;24(6):1523-7.
319. Zou W, Croke M, Fukunaga T, Broekelmann TJ, Mecham RP, Teitelbaum SL. Zap70 inhibits Syk-mediated osteoclast function. *J Cell Biochem* 2013 Aug;114(8):1871-8.

320. Borowski P, Heiland M, Kornetzky L, Medem S, Laufs R. Purification of catalytic domain of rat spleen p72syk kinase and its phosphorylation and activation by protein kinase C. *Biochem J*1998 Apr 15;331 (Pt 2):649-57.
321. Furlong MT, Mahrenholz AM, Kim KH, Ashendel CL, Harrison ML, Geahlen RL. Identification of the major sites of autophosphorylation of the murine protein-tyrosine kinase Syk. *Biochim Biophys Acta*1997 Feb 4;1355(2):177-90.
322. Wu NL, Huang DY, Tsou HN, Lin YC, Lin WW. Syk mediates IL-17-induced CCL20 expression by targeting Act1-dependent K63-linked ubiquitination of TRAF6. *J Invest Dermatol*2015 Feb;135(2):490-8.
323. Katkere B, Rosa S, Drake JR. The Syk-binding ubiquitin ligase c-Cbl mediates signaling-dependent B cell receptor ubiquitination and B cell receptor-mediated antigen processing and presentation. *J Biol Chem*2012 May 11;287(20):16636-44.
324. Hatton O, Lambert SL, Krams SM, Martinez OM. Src kinase and Syk activation initiate PI3K signaling by a chimeric latent membrane protein 1 in Epstein-Barr virus (EBV)+ B cell lymphomas. *PLoS One*2012;7(8):e42610.
325. Galan JA, Paris LL, Zhang HJ, Adler J, Geahlen RL, Tao WA. Proteomic studies of Syk-interacting proteins using a novel amine-specific isotope tag and GFP nanotrap. *J Am Soc Mass Spectrom*2011 Feb;22(2):319-28.
326. Bonilla FA, Fujita RM, Pivniouk VI, Chan AC, Geha RS. Adapter proteins SLP-76 and BLNK both are expressed by murine macrophages and are linked to signaling via Fcgamma receptors I and II/III. *Proc Natl Acad Sci U S A*2000 Feb 15;97(4):1725-30.
327. Shih CH, Chiang TB, Wang WJ. A critical role for the regulation of Syk from agglutination to aggregation in human platelets. *Biochem Biophys Res Commun*2014 Jan 10;443(2):580-5.
328. Buitrago L, Bhavanasi D, Dangelmaier C, Manne BK, Badolia R, Borgognone A, Tsygankov AY, McKenzie SE, Kunapuli SP. Tyrosine phosphorylation on spleen tyrosine kinase (Syk) is differentially regulated in human and murine platelets by protein kinase C isoforms. *J Biol Chem*2013 Oct 4;288(40):29160-9.
329. de Castro RO, Zhang J, Groves JR, Barbu EA, Siraganian RP. Once phosphorylated, tyrosines in carboxyl terminus of protein-tyrosine kinase Syk interact with signaling proteins, including TULA-2, a negative regulator of mast cell degranulation. *J Biol Chem*2012 Mar 9;287(11):8194-204.
330. Fernandez JA, Keshvara LM, Peters JD, Furlong MT, Harrison ML, Geahlen RL. Phosphorylation- and activation-independent association of the tyrosine kinase Syk and the tyrosine kinase substrates Cbl and Vav with tubulin in B-cells. *J Biol Chem*1999 Jan 15;274(3):1401-6.
331. Visperas PR, Winger JA, Horton TM, Shah NH, Aum DJ, Tao A, Barros T, Yan Q, Wilson CG, Arkin MR, Weiss A, Kuriyan J. Modification by covalent reaction or oxidation of cysteine residues in the tandem-SH2 domains of ZAP-70 and Syk can block phosphopeptide binding. *Biochem J*2015 Jan 1;465(1):149-61.
332. Hobeika E, Levit-Zerdoun E, Anastasopoulou V, Pohlmeier R, Altmeier S, Alsadeq A, Dobenecker MW, Pelanda R, Reth M. CD19 and BAFF-R can signal to promote B-cell survival in the absence of Syk. *EMBO J*2015 Jan 28.
333. Kremlitzka M, Macsik-Valent B, Erdei A. Syk is indispensable for CpG-induced activation and differentiation of human B cells. *Cell Mol Life Sci*2014 Dec 28.

334. Infantino S, Tarlinton DM. Feeling a little SYK after mixing BAFF with BCR. *Immunity* 2013 Mar 21;38(3):406-8.
335. Wang YH, Fan L, Wang L, Zhang R, Zou ZJ, Fang C, Zhang LN, Li JY, Xu W. Expression levels of Lyn, Syk, PLCgamma2 and ERK in patients with chronic lymphocytic leukemia, and higher levels of Lyn are associated with a shorter treatment-free survival. *Leuk Lymphoma* 2013 Jun;54(6):1165-70.
336. Cheng S, Coffey G, Zhang XH, Shakhovich R, Song Z, Lu P, Pandey A, Melnick AM, Sinha U, Wang YL. SYK inhibition and response prediction in diffuse large B-cell lymphoma. *Blood* 2011 Dec 8;118(24):6342-52.
337. Baert K, Sonck E, Goddeeris BM, Devriendt B, Cox E. Cell type-specific differences in beta-glucan recognition and signalling in porcine innate immune cells. *Dev Comp Immunol* 2015 Jan;48(1):192-203.
338. Uckun FM, Ma H, Ozer Z, Goodman P, Zhang J, Qazi S. A Previously Unknown Unique Challenge for Inhibitors of Syk Atp-Binding Site: Role of Syk as a Cell Cycle Checkpoint Regulator. *EBioMedicine* 2014 Nov 1;1(1):16-28.
339. Ruzza P, Biondi B, Calderan A. Therapeutic prospect of Syk inhibitors. *Expert Opin Ther Pat* 2009 Oct;19(10):1361-76.
340. Shih A, Nazi I, Kelton JG, Arnold DM. Novel treatments for immune thrombocytopenia. *Presse Med* 2014 Apr;43(4 Pt 2):e87-95.
341. Lorenz V, Stegner D, Stritt S, Vogtle T, Kiefer F, Witke W, Schymeinsky J, Watson SP, Walzog B, Nieswandt B. Targeted downregulation of platelet CLEC-2 occurs through Syk-independent internalization. *Blood* 2015 Mar 20.
342. Lu Y, Li X, Park YN, Kwon O, Piao D, Chang YC, Kim CH, Lee E, Son JK, Chang HW. Britanin Suppresses IgE/Ag-Induced Mast Cell Activation by Inhibiting the Syk Pathway. *Biomol Ther (Seoul)* 2014 May;22(3):193-9.
343. Macglashan D, Jr., Moore G, Muchhal U. Regulation of IgE-mediated signalling in human basophils by CD32b and its role in Syk down-regulation: basic mechanisms in allergic disease. *Clin Exp Allergy* 2014;44(5):713-23.
344. Harrison C. Anticancer drugs: Hitting SYK in B-ALL leukaemia. *Nat Rev Drug Discov* 2014 Jul;13(7):494.
345. Chihara K, Kimura Y, Honjo C, Takeuchi K, Sada K. Syk inhibitors. *Nihon Rinsho Meneki Gakkai Kaishi* 2013;36(4):197-202.
346. Schwartz M, Kipnis J, Rivest S, Prat A. How do immune cells support and shape the brain in health, disease, and aging? *J Neurosci* 2013 Nov 6;33(45):17587-96.
347. Diegelmann RF, Cohen IK, Kaplan AM. The role of macrophages in wound repair: a review. *Plast Reconstr Surg* 1981 Jul;68(1):107-13.
348. LeFevre ME, Hammer R, Joel DD. Macrophages of the mammalian small intestine: a review. *J Reticuloendothel Soc* 1979 Nov;26(5):553-73.
349. Leder LD. The origin of blood monocytes and macrophages. A review. *Blut* 1967 Nov;16(2):86-98.
350. Lin TJ, Hirji N, Stenton GR, Gilchrist M, Grill BJ, Schreiber AD, Befus AD. Activation of macrophage CD8: pharmacological studies of TNF and IL-1 beta production. *J Immunol* 2000 Feb 15;164(4):1783-92.

351. Hirji NS, Lin TJ, Gilchrist M, Nault G, Nohara O, Grill BJ, Belosevic M, Stenton GR, Schreiber AD, Befus AD. Novel CD8 molecule on macrophages and mast cells: expression, function and signaling. *Int Arch Allergy Immunol* 1999 Feb-Apr;118(2-4):180-2.
352. Yang WS, Jeong D, Yi YS, Lee BH, Kim TW, Htwe KM, Kim YD, Yoon KD, Hong S, Lee WS, Cho JY. Myrsine seguinii ethanolic extract and its active component quercetin inhibit macrophage activation and peritonitis induced by LPS by targeting to Syk/Src/IRAK-1. *J Ethnopharmacol* 2014 Feb 12;151(3):1165-74.
353. Aronoff DM, Serezani CH, Carstens JK, Marshall T, Gangireddy SR, Peters-Golden M, Reddy RC. Stimulatory Effects of Peroxisome Proliferator-Activated Receptor-gamma on Fcgamma Receptor-Mediated Phagocytosis by Alveolar Macrophages. *PPAR Res* 2007;2007:52546.
354. Jaumouille V, Farkash Y, Jaqaman K, Das R, Lowell CA, Grinstein S. Actin cytoskeleton reorganization by Syk regulates Fcgamma receptor responsiveness by increasing its lateral mobility and clustering. *Dev Cell* 2014 Jun 9;29(5):534-46.
355. Roy R, Chauhan LK, Das M, Tripathi A, Dwivedi PD. Phagocytic cells internalize ZnO particles by FcgammaII/III-receptor pathway. *Immunobiology* 2014 Oct;219(10):746-55.
356. Greuber EK, Pendergast AM. Abl family kinases regulate FcgammaR-mediated phagocytosis in murine macrophages. *J Immunol* 2012 Dec 1;189(11):5382-92.
357. Fang J, Wang Y, Lv X, Shen X, Ni X, Ding K. Structure of a beta-glucan from *Grifola frondosa* and its antitumor effect by activating Dectin-1/Syk/NF-kappaB signaling. *Glycoconj J* 2012 Aug;29(5-6):365-77.
358. Ientile R, Curro M, Caccamo D. Transglutaminase 2 and neuroinflammation. *Amino Acids* 2015 Jan;47(1):19-26.
359. Okamoto F, Saeki K, Sumimoto H, Yamasaki S, Yokomizo T. Leukotriene B4 augments and restores Fc gammaRs-dependent phagocytosis in macrophages. *J Biol Chem* 2010 Dec 24;285(52):41113-21.
360. Nakamura M, Watanabe N. Ubiquitin-like protein MNSFbeta/endophilin II complex regulates Dectin-1-mediated phagocytosis and inflammatory responses in macrophages. *Biochem Biophys Res Commun* 2010 Oct 15;401(2):257-61.
361. Cash JL, Christian AR, Greaves DR. Chemerin peptides promote phagocytosis in a ChemR23- and Syk-dependent manner. *J Immunol* 2010 May 1;184(9):5315-24.
362. Sato N, Kim MK, Schreiber AD. Enhancement of fcgamma receptor-mediated phagocytosis by transforming mutants of Cbl. *J Immunol* 1999 Dec 1;163(11):6123-31.
363. Larbolette O, Wollscheid B, Schweikert J, Nielsen PJ, Wienands J. SH3P7 is a cytoskeleton adapter protein and is coupled to signal transduction from lymphocyte antigen receptors. *Mol Cell Biol* 1999 Feb;19(2):1539-46.
364. Indik ZK, Park JG, Hunter S, Schreiber AD. The molecular dissection of Fc gamma receptor mediated phagocytosis. *Blood* 1995 Dec 15;86(12):4389-99.
365. Dung TT, Kim SC, Yoo BC, Sung GH, Yang WS, Kim HG, Park JG, Rhee MH, Park KW, Yoon K, Lee Y, Hong S, Kim JH, Cho JY. (5-Hydroxy-4-oxo-4H-pyran-2-yl)methyl 6-hydroxynaphthalene-2-carboxylate, a kojic acid derivative, inhibits inflammatory mediator production via the suppression of Syk/Src and NF-kappaB activation. *Int Immunopharmacol* 2014 May;20(1):37-45.

366. Kim E, Yoon KD, Lee WS, Yang WS, Kim SH, Sung NY, Baek KS, Kim Y, Htwe KM, Kim YD, Hong S, Kim JH, Cho JY. Syk/Src-targeted anti-inflammatory activity of *Codariocalyx motorius* ethanolic extract. *J Ethnopharmacol*2014 Aug 8;155(1):185-93.
367. Miller YI, Choi SH, Wiesner P, Bae YS. The SYK side of TLR4: signalling mechanisms in response to LPS and minimally oxidized LDL. *Br J Pharmacol*2012 Nov;167(5):990-9.
368. Kankkunen P, Teirila L, Rintahaka J, Alenius H, Wolff H, Matikainen S. (1,3)-beta-glucans activate both dectin-1 and NLRP3 inflammasome in human macrophages. *J Immunol*2010 Jun 1;184(11):6335-42.
369. Zhang S, Yang L, Kouadir M, Tan R, Lu Y, Chang J, Xu B, Yin X, Zhou X, Zhao D. PP2 and piceatannol inhibit PrP106-126-induced iNOS activation mediated by CD36 in BV2 microglia. *Acta Biochim Biophys Sin (Shanghai)*2013 Sep;45(9):763-72.
370. Takamiya R, Ohtsubo K, Takamatsu S, Taniguchi N, Angata T. The interaction between Siglec-15 and tumor-associated sialyl-Tn antigen enhances TGF-beta secretion from monocytes/macrophages through the DAP12-Syk pathway. *Glycobiology*2013 Feb;23(2):178-87.
371. Hughes CE, Finney BA, Koentgen F, Lowe KL, Watson SP. The N-terminal SH2 domain of Syk is required for (hem)ITAM, but not integrin, signaling in mouse platelets. *Blood*2015 Jan 1;125(1):144-54.
372. Falker K, Klarstrom-Engstrom K, Bengtsson T, Lindahl TL, Grenegard M. The toll-like receptor 2/1 (TLR2/1) complex initiates human platelet activation via the src/Syk/LAT/PLCgamma2 signalling cascade. *Cell Signal*2014 Feb;26(2):279-86.
373. Hatterer E, Benon A, Chounlamountri N, Watrin C, Angibaud J, Jouanneau E, Boudin H, Honnorat J, Pellier-Monnin V, Noraz N. Syk kinase is phosphorylated in specific areas of the developing nervous system. *Neurosci Res*2011 Jun;70(2):172-82.
374. Armstrong RC, Mierzwa AJ, Sullivan GM, Sanchez MA. Myelin and oligodendrocyte lineage cells in white matter pathology and plasticity after traumatic brain injury. *Neuropharmacology*2015 May 9.
375. Louveau A, Smirnov I, Keyes TJ, Eccles JD, Rouhani SJ, Peske JD, Derecki NC, Castle D, Mandell JW, Lee KS, Harris TH, Kipnis J. Structural and functional features of central nervous system lymphatic vessels. *Nature*2015 Jul 16;523(7560):337-41.
376. Combs CK, Karlo JC, Kao SC, Landreth GE. beta-Amyloid stimulation of microglia and monocytes results in TNFalpha-dependent expression of inducible nitric oxide synthase and neuronal apoptosis. *J Neurosci*2001 Feb 15;21(4):1179-88.
377. Baudouin SJ, Angibaud J, Loussouarn G, Bonnamain V, Matsuura A, Kinebuchi M, Naveilhan P, Boudin H. The signaling adaptor protein CD3zeta is a negative regulator of dendrite development in young neurons. *Mol Biol Cell*2008 Jun;19(6):2444-56.
378. Tsujimura T, Yanagi S, Inatome R, Takano T, Ishihara I, Mitsui N, Takahashi S, Yamamura H. Syk protein-tyrosine kinase is involved in neuron-like differentiation of embryonal carcinoma P19 cells. *FEBS Lett*2001 Feb 2;489(2-3):129-33.
379. He X, Sun BG. [Roles of microglia and immune receptors in Alzheimer's disease]. *Yao Xue Xue Bao*2014 Jun;49(6):774-80.

380. McDonald DR, Brunden KR, Landreth GE. Amyloid fibrils activate tyrosine kinase-dependent signaling and superoxide production in microglia. *J Neurosci*1997 Apr 1;17(7):2284-94.
381. Mattson MP, Rydel RE. Alzheimer's disease. Amyloid ox-tox transducers. *Nature*1996 Aug 22;382(6593):674-5.
382. Neumann H, Kotter MR, Franklin RJ. Debris clearance by microglia: an essential link between degeneration and regeneration. *Brain*2009 Feb;132(Pt 2):288-95.
383. Dietrich J, Cella M, Seiffert M, Buhring HJ, Colonna M. Cutting edge: signal-regulatory protein beta 1 is a DAP12-associated activating receptor expressed in myeloid cells. *J Immunol*2000 Jan 1;164(1):9-12.
384. Jin SC, Benitez BA, Karch CM, Cooper B, Skorupa T, Carrell D, Norton JB, Hsu S, Harari O, Cai Y, Bertelsen S, Goate AM, Cruchaga C. Coding variants in TREM2 increase risk for Alzheimer's disease. *Hum Mol Genet*2014 Nov 1;23(21):5838-46.
385. Linnartz-Gerlach B, Mathews M, Neumann H. Sensing the neuronal glycocalyx by glial sialic acid binding immunoglobulin-like lectins. *Neuroscience*2014 Sep 5;275:113-24.
386. Angata T, Kerr SC, Greaves DR, Varki NM, Crocker PR, Varki A. Cloning and characterization of human Siglec-11. A recently evolved signaling molecule that can interact with SHP-1 and SHP-2 and is expressed by tissue macrophages, including brain microglia. *J Biol Chem*2002 Jul 5;277(27):24466-74.
387. Blasi E, Barluzzi R, Bocchini V, Mazzolla R, Bistoni F. Immortalization of murine microglial cells by a v-raf/v-myc carrying retrovirus. *J Neuroimmunol*1990 May;27(2-3):229-37.
388. Lourenco MV, Ledo JH. Targeting Alzheimer's pathology through PPARgamma signaling: modulation of microglial function. *J Neurosci*2013 Mar 20;33(12):5083-4.
389. Xu S, Guan Q, Wang C, Wei X, Chen X, Zheng B, An P, Zhang J, Chang L, Zhou W, Mody I, Wang Q. Rosiglitazone prevents the memory deficits induced by amyloid-beta oligomers via inhibition of inflammatory responses. *Neurosci Lett*2014 Aug 22;578:7-11.
390. Kedersha N, Anderson P. Stress granules: sites of mRNA triage that regulate mRNA stability and translatability. *Biochem Soc Trans*2002 Nov;30(Pt 6):963-9.
391. Kedersha N, Ivanov P, Anderson P. Stress granules and cell signaling: more than just a passing phase? *Trends Biochem Sci*2013 Oct;38(10):494-506.
392. Anderson P, Kedersha N, Ivanov P. Stress granules, P-bodies and cancer. *Biochim Biophys Acta*2014 Dec 5.
393. Kedersha N, Cho MR, Li W, Yacono PW, Chen S, Gilks N, Golan DE, Anderson P. Dynamic shuttling of TIA-1 accompanies the recruitment of mRNA to mammalian stress granules. *J Cell Biol*2000 Dec 11;151(6):1257-68.
394. Kedersha N, Anderson P. Mammalian stress granules and processing bodies. *Methods Enzymol*2007;431:61-81.
395. Hofmann I, Casella M, Schnolzer M, Schlechter T, Spring H, Franke WW. Identification of the junctional plaque protein plakophilin 3 in cytoplasmic particles containing RNA-binding proteins and the recruitment of plakophilins 1 and 3 to stress granules. *Mol Biol Cell*2006 Mar;17(3):1388-98.

396. Kedersha NL, Gupta M, Li W, Miller I, Anderson P. RNA-binding proteins TIA-1 and TIAR link the phosphorylation of eIF-2 alpha to the assembly of mammalian stress granules. *J Cell Biol*1999 Dec 27;147(7):1431-42.
397. Shelkownikova TA, Robinson HK, Connor-Robson N, Buchman VL. Recruitment into stress granules prevents irreversible aggregation of FUS protein mislocalized to the cytoplasm. *Cell Cycle*2013 Oct 1;12(19):3194-202.
398. Bley N, Lederer M, Pfalz B, Reinke C, Fuchs T, Glass M, Moller B, Huttelmaier S. Stress granules are dispensable for mRNA stabilization during cellular stress. *Nucleic Acids Res*2015 Feb 27;43(4):e26.
399. Decker CJ, Parker R. P-bodies and stress granules: possible roles in the control of translation and mRNA degradation. *Cold Spring Harb Perspect Biol*2012 Sep;4(9):a012286.
400. Martin S, Tazi J. Visualization of G3BP stress granules dynamics in live primary cells. *J Vis Exp*2014(87).
401. Ohn T, Anderson P. The role of posttranslational modifications in the assembly of stress granules. *Wiley Interdiscip Rev RNA*2010 Nov-Dec;1(3):486-93.
402. Souquere S, Mollet S, Kress M, Dautry F, Pierron G, Weil D. Unravelling the ultrastructure of stress granules and associated P-bodies in human cells. *J Cell Sci*2009 Oct 15;122(Pt 20):3619-26.
403. Tourriere H, Chebli K, Zekri L, Courselaud B, Blanchard JM, Bertrand E, Tazi J. The RasGAP-associated endoribonuclease G3BP assembles stress granules. *J Cell Biol*2003 Mar 17;160(6):823-31.
404. Kedersha N, Stoecklin G, Ayodele M, Yacono P, Lykke-Andersen J, Fritzler MJ, Scheuner D, Kaufman RJ, Golan DE, Anderson P. Stress granules and processing bodies are dynamically linked sites of mRNP remodeling. *J Cell Biol*2005 Jun 20;169(6):871-84.
405. Matus S, Bosco DA, Hetz C. Autophagy meets fused in sarcoma-positive stress granules. *Neurobiol Aging*2014 Dec;35(12):2832-5.
406. Li YR, King OD, Shorter J, Gitler AD. Stress granules as crucibles of ALS pathogenesis. *J Cell Biol*2013 Apr 29;201(3):361-72.
407. Anderson P, Kedersha N. Visibly stressed: the role of eIF2, TIA-1, and stress granules in protein translation. *Cell Stress Chaperones*2002 Apr;7(2):213-21.
408. Abrahamyan LG, Chatel-Chaix L, Ajamian L, Milev MP, Monette A, Clement JF, Song R, Lehmann M, DesGroseillers L, Laughrea M, Boccaccio G, Mouland AJ. Novel Staufen1 ribonucleoproteins prevent formation of stress granules but favour encapsidation of HIV-1 genomic RNA. *J Cell Sci*2010 Feb 1;123(Pt 3):369-83.
409. Tsai NP, Ho PC, Wei LN. Regulation of stress granule dynamics by Grb7 and FAK signalling pathway. *EMBO J*2008 Mar 5;27(5):715-26.
410. Vanderweyde T, Youmans K, Liu-Yesucevitz L, Wolozin B. Role of stress granules and RNA-binding proteins in neurodegeneration: a mini-review. *Gerontology*2013;59(6):524-33.
411. Solomon S, Xu Y, Wang B, David MD, Schubert P, Kennedy D, Schrader JW. Distinct structural features of caprin-1 mediate its interaction with G3BP-1 and its induction of phosphorylation of eukaryotic translation initiation factor 2alpha, entry to cytoplasmic stress granules, and selective interaction with a subset of mRNAs. *Mol Cell Biol*2007 Mar;27(6):2324-42.

412. Hua Y, Zhou J. Survival motor neuron protein facilitates assembly of stress granules. *FEBS Lett*2004 Aug 13;572(1-3):69-74.
413. Kim WJ, Back SH, Kim V, Ryu I, Jang SK. Sequestration of TRAF2 into stress granules interrupts tumor necrosis factor signaling under stress conditions. *Mol Cell Biol*2005 Mar;25(6):2450-62.
414. Moutaoufik MT, El Fatimy R, Nassour H, Gareau C, Lang J, Tanguay RM, Mazroui R, Khandjian EW. UVC-induced stress granules in mammalian cells. *PLoS One*2014;9(11):e112742.
415. Bounedjah O, Desforges B, Wu TD, Pioche-Durieu C, Marco S, Hamon L, Curmi PA, Guerquin-Kern JL, Pietrement O, Pastre D. Free mRNA in excess upon polysome dissociation is a scaffold for protein multimerization to form stress granules. *Nucleic Acids Res*2014 Jul;42(13):8678-91.
416. Yang WH, Yu JH, Gulick T, Bloch KD, Bloch DB. RNA-associated protein 55 (RAP55) localizes to mRNA processing bodies and stress granules. *RNA*2006 Apr;12(4):547-54.
417. Arimoto K, Fukuda H, Imajoh-Ohmi S, Saito H, Takekawa M. Formation of stress granules inhibits apoptosis by suppressing stress-responsive MAPK pathways. *Nat Cell Biol*2008 Nov;10(11):1324-32.
418. Kaehler C, Isensee J, Hucho T, Lehrach H, Krobisch S. 5-Fluorouracil affects assembly of stress granules based on RNA incorporation. *Nucleic Acids Res*2014 Jun;42(10):6436-47.
419. Wolozin B. Physiological protein aggregation run amuck: stress granules and the genesis of neurodegenerative disease. *Discov Med*2014 Jan;17(91):47-52.
420. Vanderweyde T, Yu H, Varnum M, Liu-Yesucevitz L, Citro A, Ikezu T, Duff K, Wolozin B. Contrasting pathology of the stress granule proteins TIA-1 and G3BP in tauopathies. *J Neurosci*2012 Jun 13;32(24):8270-83.
421. Reineke LC, Kedersha N, Langereis MA, van Kuppeveld FJ, Lloyd RE. Stress Granules Regulate Double-Stranded RNA-Dependent Protein Kinase Activation through a Complex Containing G3BP1 and Caprin1. *MBio*2015;6(2).
422. Figley MD, Bieri G, Kolaitis RM, Taylor JP, Gitler AD. Profilin 1 associates with stress granules and ALS-linked mutations alter stress granule dynamics. *J Neurosci*2014 Jun 11;34(24):8083-97.
423. Dewey CM, Cenik B, Sephton CF, Johnson BA, Herz J, Yu G. TDP-43 aggregation in neurodegeneration: are stress granules the key? *Brain Res*2012 Jun 26;1462:16-25.
424. Zou T, Yang X, Pan D, Huang J, Sahin M, Zhou J. SMN deficiency reduces cellular ability to form stress granules, sensitizing cells to stress. *Cell Mol Neurobiol*2011 May;31(4):541-50.
425. Hua Y, Zhou J. Rpp20 interacts with SMN and is re-distributed into SMN granules in response to stress. *Biochem Biophys Res Commun*2004 Jan 30;314(1):268-76.
426. Righi M, Mori L, De Libero G, Sironi M, Biondi A, Mantovani A, Donini SD, Ricciardi-Castagnoli P. Monokine production by microglial cell clones. *Eur J Immunol*1989 Aug;19(8):1443-8.

427. Wang WH, Childress MO, Geahlen RL. Syk interacts with and phosphorylates nucleolin to stabilize Bcl-x(L) mRNA and promote cell survival. *Mol Cell Biol*2014 Oct;34(20):3788-99.
428. Moussaud S, Draheim HJ. A new method to isolate microglia from adult mice and culture them for an extended period of time. *J Neurosci Methods*2010 Mar 30;187(2):243-53.
429. Lee JK, Tansey MG. Microglia isolation from adult mouse brain. *Methods Mol Biol*2013;1041:17-23.
430. Pan XD, Zhu YG, Lin N, Zhang J, Ye QY, Huang HP, Chen XC. Microglial phagocytosis induced by fibrillar beta-amyloid is attenuated by oligomeric beta-amyloid: implications for Alzheimer's disease. *Mol Neurodegener*2011;6:45.
431. Fiala M, Liu PT, Espinosa-Jeffrey A, Rosenthal MJ, Bernard G, Ringman JM, Sayre J, Zhang L, Zaghi J, Dejbakhsh S, Chiang B, Hui J, Mahanian M, Baghaee A, Hong P, Cashman J. Innate immunity and transcription of MGAT-III and Toll-like receptors in Alzheimer's disease patients are improved by bisdemethoxycurcumin. *Proc Natl Acad Sci U S A*2007 Jul 31;104(31):12849-54.
432. Khanna R, Roy L, Zhu X, Schlichter LC. K⁺ channels and the microglial respiratory burst. *Am J Physiol Cell Physiol*2001 Apr;280(4):C796-806.
433. Uy B, McGlashan SR, Shaikh SB. Measurement of reactive oxygen species in the culture media using Acridan Lumigen PS-3 assay. *J Biomol Tech*2011 Sep;22(3):95-107.
434. McEwen E, Kedersha N, Song B, Scheuner D, Gilks N, Han A, Chen JJ, Anderson P, Kaufman RJ. Heme-regulated inhibitor kinase-mediated phosphorylation of eukaryotic translation initiation factor 2 inhibits translation, induces stress granule formation, and mediates survival upon arsenite exposure. *J Biol Chem*2005 Apr 29;280(17):16925-33.
435. Yamaguchi A, Kitajo K. The effect of PRMT1-mediated arginine methylation on the subcellular localization, stress granules, and detergent-insoluble aggregates of FUS/TLS. *PLoS One*2012;7(11):e49267.
436. Parker SJ, Meyerowitz J, James JL, Liddell JR, Crouch PJ, Kanninen KM, White AR. Endogenous TDP-43 localized to stress granules can subsequently form protein aggregates. *Neurochem Int*2012 Mar;60(4):415-24.
437. Wolozin B. Cholesterol and the biology of Alzheimer's disease. *Neuron*2004 Jan 8;41(1):7-10.
438. Cameron B, Landreth GE. Inflammation, microglia, and Alzheimer's disease. *Neurobiol Dis*2010 Mar;37(3):503-9.
439. Mocsai A, Abram CL, Jakus Z, Hu Y, Lanier LL, Lowell CA. Integrin signaling in neutrophils and macrophages uses adaptors containing immunoreceptor tyrosine-based activation motifs. *Nat Immunol*2006 Dec;7(12):1326-33.
440. Vlad SC, Miller DR, Kowall NW, Felson DT. Protective effects of NSAIDs on the development of Alzheimer disease. *Neurology*2008 May 6;70(19):1672-7.
441. Heit B, Kim H, Cosio G, Castano D, Collins R, Lowell CA, Kain KC, Trimble WS, Grinstein S. Multimolecular signaling complexes enable Syk-mediated signaling of CD36 internalization. *Dev Cell*2013 Feb 25;24(4):372-83.

442. Bamberger ME, Harris ME, McDonald DR, Husemann J, Landreth GE. A cell surface receptor complex for fibrillar beta-amyloid mediates microglial activation. *J Neurosci*2003 Apr 1;23(7):2665-74.
443. Schieven GL, Kirihaara JM, Burg DL, Geahlen RL, Ledbetter JA. p72syk tyrosine kinase is activated by oxidizing conditions that induce lymphocyte tyrosine phosphorylation and Ca²⁺ signals. *J Biol Chem*1993 Aug 5;268(22):16688-92.
444. Dodel RC, Du Y, Depboylu C, Hampel H, Frolich L, Haag A, Hemmeter U, Paulsen S, Teipel SJ, Brettschneider S, Spottke A, Nolker C, Moller HJ, Wei X, Farlow M, Sommer N, Oertel WH. Intravenous immunoglobulins containing antibodies against beta-amyloid for the treatment of Alzheimer's disease. *J Neurol Neurosurg Psychiatry*2004 Oct;75(10):1472-4.
445. Relkin NR, Szabo P, Adamiak B, Burgut T, Monthe C, Lent RW, Younkin S, Younkin L, Schiff R, Weksler ME. 18-Month study of intravenous immunoglobulin for treatment of mild Alzheimer disease. *Neurobiol Aging*2009 Nov;30(11):1728-36.
446. Loeffler DA. Intravenous immunoglobulin and Alzheimer's disease: what now? *J Neuroinflammation*2013;10:70.
447. Sudduth TL, Greenstein A, Wilcock DM. Intracranial injection of Gammagard, a human IVIg, modulates the inflammatory response of the brain and lowers Abeta in APP/PS1 mice along a different time course than anti-Abeta antibodies. *J Neurosci*2013 Jun 5;33(23):9684-92.
448. Puig KL, Lutz BM, Urquhart SA, Rebel AA, Zhou X, Manocha GD, Sens M, Tuteja AK, Foster NL, Combs CK. Overexpression of Mutant Amyloid-beta Protein Precursor and Presenilin 1 Modulates Enteric Nervous System. *J Alzheimers Dis*2015 Jan 1;44(4):1263-78.
449. Fu Y, Hsiao JH, Paxinos G, Halliday GM, Kim WS. ABCA5 regulates amyloid-beta peptide production and is associated with Alzheimer's disease neuropathology. *J Alzheimers Dis*2015;43(3):857-69.
450. Zeng KW, Wang S, Dong X, Jiang Y, Jin HW, Tu PF. Sesquiterpene dimmer (DSF-27) inhibits the release of neuroinflammatory mediators from microglia by targeting spleen tyrosine kinase (Syk) and Janus kinase 2 (Jak2): Two major non-receptor tyrosine signaling proteins involved in inflammatory events. *Toxicol Appl Pharmacol*2014 Mar 15;275(3):244-56.
451. Fanger NA, Voigtlaender D, Liu C, Swink S, Wardwell K, Fisher J, Graziano RF, Pfefferkorn LC, Guyre PM. Characterization of expression, cytokine regulation, and effector function of the high affinity IgG receptor Fc gamma RI (CD64) expressed on human blood dendritic cells. *J Immunol*1997 Apr 1;158(7):3090-8.
452. Oh H, Ozkirimli E, Shah K, Harrison ML, Geahlen RL. Generation of an analog-sensitive Syk tyrosine kinase for the study of signaling dynamics from the B cell antigen receptor. *J Biol Chem*2007 Nov 16;282(46):33760-8.

VITA

VITA

Soumitra Ghosh was born on April 20, 1987 in Mumbai, India. He moved to Kolkata soon after and finished his schooling at Kendriya Vidyalaya Cossipore. Right from childhood he dreamt about becoming a scientist. Among all the science subjects he learned as a kid, biology appealed him the most. To fulfill his dreams he decided to major in Genetic Engineering as an undergraduate. Four years later in 2009, Soumitra completed his Bachelor of Technology degree in Genetic Engineering from SRM University, Chennai, India. During his undergraduate to obtain research experience, he spent all his summer vacation as an intern in various Universities and national labs. In his junior year, he managed to get a summer fellowship from Department of Biotechnology, India to pursue neuroscience research at ACBR, Delhi. He soon realized that neuroscience was his area of interest and decided to obtain a PhD. in neuroscience. In August 2009 right after graduation, he joined Purdue University as a graduate student to pursue his neuroscience research. Initially he worked in Department of Chemistry and later moved to Geahlen lab. Under excellent guidance and support of Dr. Robert Geahlen, he studied the role of Syk in microglial cell and stress granules. In the fall of 2015 he completed his PhD in Medicinal Chemistry and Molecular Pharmacology with a concentration in integrative neuroscience. He is going to continue as a post doctoral researcher in neuroimmunology and chase his dreams of becoming an independent scientist.

PUBLICATIONS

PUBLICATIONS

Ghosh S and Geahlen LR. Stress Granules modulate Syk to cause Microglial Dysfunction in Alzheimer's disease. 2015 (EBioMedicine , In press, September 2015)

Krisenko MO, Higgins RL, Ghosh S Zhou Q, Trybula JS, Wang WH, and Geahlen RL. Syk is Recruited to Stress Granules and Promotes their Clearance through Autophagy. (The Journal of Biological Chemistry, In press, October 2015)

Johnson EO, Chang KH, Ghosh S, Venkatesh C, Giger K, Low PS and Shah K. LIMK2 is a crucial regulator and effector of Aurora-A-kinase-mediated malignancy. Journal of Cell Science. 2012 Mar 1;125(Pt 5):1204-16.

Kumar D, Maruthi Kumar N, Ghosh S, Shah K. Novel bis(indolyl) hydrazide-hydrazones as potent cytotoxic agents. Bioorg Med Chem Lett. 2012 Jan 1; 22(1):212-5.

Chang KH, Multani PS, Sun KH, Vincent F, Pablo Y, Ghosh S, Gupta R, Lee HP, Lee HG, Smith MA and Shah K. Nuclear envelope dispersion triggered by deregulated Cdk5 precedes neuronal death. Molecular Biology of the Cell. 2011 May; 22(9):1452-62.

Sun KH, Chang KH, Ghosh S, Clawson S, Mirzaei H, Regnier F, Shah K. GSTP1 is a Critical Regulator of Cdk5 Kinase Activity in Alzheimer's disease. Journal of Neurochemistry. 2011 Sep; 118(5):902-14.

Johnson EO, Chang KH, de Pablo Y, Ghosh S, Mehta R, Badve S, Shah K. PHLDA1 is a crucial negative regulator and effector of Aurora A kinase in breast cancer. *Journal of Cell Science*. 2011 Aug 15;124(Pt 16):2711-22.